

Biased Oxytocinergic Modulation of Midbrain Dopamine Systems

Highlights

- PVN oxytocinergic neurons project to midbrain dopamine regions
- VTA and SNc express multiple oxytocin-sensing GPCRs
- Oxytocin release enhances the activity of VTA DA neurons
- Oxytocin release decreases the activity of SNc DA neurons via inhibitory circuits

Authors

Lei Xiao, Michael F. Priest,
Jordan Nasenbeny, Ting Lu,
Yevgenia Kozorovitskiy

Correspondence

yevgenia.kozorovitskiy@
northwestern.edu

In Brief

Xiao et al. reveal oxytocinergic projections from PVN to VTA and SNc, where, by different circuit and receptor mechanisms, oxytocin release biases dopamine neuron activity in opposite directions. This work places dopamine signaling under prominent control of a peptide neuromodulator.



Biased Oxytocinergic Modulation of Midbrain Dopamine Systems

Lei Xiao,¹ Michael F. Priest,¹ Jordan Nasenbeny,¹ Ting Lu,² and Yevgenia Kozorovitskiy^{1,3,*}

¹Department of Neurobiology

²Roxelyn and Richard Pepper Department of Communication Sciences and Disorders
Northwestern University, Evanston, IL 60208, USA

³Lead Contact

*Correspondence: yevgenia.kozorovitskiy@northwestern.edu

<http://dx.doi.org/10.1016/j.neuron.2017.06.003>

SUMMARY

The release of dopamine (DA) regulates rewarding behavior and motor actions through striatum-targeting efferents from ventral tegmental area (VTA) and substantia nigra pars compacta (SNc). Here, we map and functionally characterize axonal projections from oxytocin neurons in the hypothalamic paraventricular nucleus to midbrain DA regions. Electrophysiological recordings of DA neurons reveal that both the application of oxytocin and optogenetic stimulation of oxytocinergic terminals suffice to increase DA neuron activity in the VTA but downregulate it in SNc. This biased modulation is mediated by oxytocin and vasopressin G-protein-coupled receptors. Oxytocin release directly activates DA neurons and indirectly inhibits them through local GABA neurons, but the relative magnitudes of the two mechanisms differ in VTA and SNc. Oxytocin-modulated DA neurons give rise to canonical striatal projections. Since hypothalamic oxytocinergic projections also target the striatum, oxytocin is poised to bias the balance of DA tone through multiple sites in vertebrate reward circuits.

INTRODUCTION


In the vertebrate basal ganglia, timed release of the neuromodulator dopamine (DA) supports complex, goal-directed actions (Bromberg-Martin et al., 2010; Schultz, 2007; Wise, 2004). The majority of midbrain neurons that produce DA reside in the ventral tegmental area (VTA) and substantia nigra pars compacta (SNc). The core projections of VTA DA neurons target ventral striatal (vStr) areas, subserving reward-based learning, while SNc DA neurons primarily target motor action-associated dorsal striatum (dStr). Genetically guided tracing experiments have facilitated the identification of input and output wiring of DA neurons in the VTA and SNc (Beier et al., 2015; Lerner et al., 2015; Watabe-Uchida et al., 2012). The results of functional experiments, including calcium imaging of DA terminals, further support the preferential links of VTA DA signaling to

reward and SNc-derived DA tone to locomotion (Howe and Dombeck, 2016).

Oxytocin (Oxt) is synthesized and released by neurons of the hypothalamic paraventricular nucleus (PVN) and supraoptic nucleus (SON). This neuropeptide and hormone is linked to the reward system through its effects on social behaviors, including social reward, social learning, pair bond formation, parenting, and mating (Gimpl and Fahrenholz, 2001; Love, 2014; Dölen et al., 2013; Marlin et al., 2015; Choe et al., 2015). Numerous studies demonstrate an indirect or downstream interaction of Oxt and DA signaling. A key example is Oxt release into the vStr, which promotes affiliative social behavior (Liu and Wang, 2003; Shahrokh et al., 2010; Young and Wang, 2004). However, the only current evidence for a direct Oxt regulation of DA neuron activity focuses on prolactin induction in the arcuate nucleus (Briffaud et al., 2015).

Prior results suggest the possibility that Oxt also acts in the midbrain. Behavioral studies show that activation of Oxt receptors (OxtRs) in the VTA is important for social reward (Song et al., 2016). Oxt in the VTA may generally influence reward-based behavior, since Oxt infusions there suppress sucrose intake (Mullis et al., 2013). Moreover, a recent *trans*-synaptic tracing analysis showed that Oxt neurons from the PVN project to VTA DA neurons (Beier et al., 2015), while another study demonstrated that Oxt could regulate the activity of a subset of unclassified VTA neurons (Tang et al., 2014). Therefore, Oxt may directly modulate VTA DA neurons. Whether Oxt projections also target SNc DA neurons has remained unexplored. One of the two brain regions enriched in Oxt somata, the PVN, sends monosynaptic projections to the SNc (Watabe-Uchida et al., 2012), where Oxt infusions reduce locomotor activity (Angioni et al., 2016). Despite experimental support for the existence of midbrain-targeting Oxt projections and behavioral effects of Oxt infusions, the functional impact of Oxt on the activity of DA neurons in either VTA or SNc has not been examined.

In the central nervous system, Oxt binds to a single G-protein-coupled receptor (GPCR) (OxtR) primarily coupled to G_{α_q} protein effectors, but, at higher concentrations, Oxt is able to recruit G_{α_i} subunits (Busnelli et al., 2012). Binding to G_{α_q} -coupled OxtR stimulates the phospholipase C pathway to enhance neuronal firing (Gimpl and Fahrenholz, 2001; Stoop, 2012). Oxt can also bind with lower affinity to the three vasopressin (Avp) receptors V1aR, V1bR, and V2R (Tribollet et al., 1988). *Oxtr* transcripts have been found in rodent VTA and SNc (Peris et al., 2017;



(legend on next page)

Vaccari et al., 1998), while the evidence for AvpR expression in midbrain DA regions is conflicting (Dubois-Dauphin et al., 1996; Ostrowski et al., 1994; Insel et al., 1994; Tribollet et al., 1988). If these areas express both OxtR and AvpR, then the activation of either receptor by Oxt could result in a net increase in neuronal activity.

To dissect the mechanisms of oxytocinergic modulation of DA neurons in the VTA and SNc, we used a combination of anatomical, optogenetic, and electrophysiological approaches. Here, we show that Oxt controls neuronal activity in midbrain DA regions through direct and polysynaptic effects. These effects are driven by Oxt release itself, which acts on several Oxt-sensing receptors, including Oxt and Avp GPCRs. Our results place DA signaling under prominent control of a neuropeptide that is known to regulate social behavior and interact with reward signaling.

RESULTS

PVN Oxt Projections Directly Target the VTA and SNc

To determine whether Oxt neurons project to midbrain DA regions, we unilaterally injected green retrobeads (GRBs) into either the VTA or the SNc of postnatal day 25–30 (P25–P30) mice expressing tdTomato under control of the *Oxt* promoter (*Oxt*^{i-Cre}; Ai14) (Figures 1A–1F). We validated the specificity of this line by imaging the overlap between tdTomato and Oxt immunofluorescent (IF) labeling in the PVN (695/732 tdTomato⁺ neurons were Oxt⁺, 3 mice) (Figure 1A). GRB⁺ cells were observed in the ipsilateral PVN, as well as in the striatum and lateral hypothalamus (Watabe-Uchida et al., 2012), but not in the SON (Figures 1D and 1E; Figure S1A). Similar numbers of PVN neurons were labeled following GRB injections targeting the VTA or the SNc (VTA_{target}, 90.8 ± 12.2 neurons/mouse; SNc_{target}, 108.0 ± 12.5 neurons/mouse; *p* = 0.548, Mann-Whitney test, *n* = 5 mice/group) (Figures 1E and 1F). Approximately 20% of GRB⁺ neurons in the PVN were tdTomato⁺, providing evidence for a direct oxytocinergic PVN projection to midbrain DA regions (proportion tdTomato⁺/GRB⁺ in the VTA, 0.237 ± 0.022; SNc, 0.239 ± 0.016; *p* = 0.691, Mann-Whitney test, *n* = 5 mice/group) (Fig-

ure 1F). The proportion of Oxt neurons in these PVN-arising efferents is consistent with a recent *trans*-synaptic tracing analysis of PVN outputs (Beier et al., 2015).

To evaluate whether the same set of Oxt PVN neurons project to both VTA and SNc, we injected both GRBs and red retrobeads (RRBs) separately into the VTA and SNc of wild-type mice (Figure S1B). After 7–9 days, we evaluated the overlap between RB⁺ cells and Oxt neurons in the PVN (Figures 1G and 1H; Figure S1B). Both GRBs and RRBs were found in PVN Oxt neurons, at similar proportions as for single-color RB labeling. However, few GRB⁺/RRB⁺ neurons overlapped with Oxt neurons, demonstrating that Oxt projections to the VTA and SNc are largely separate (25.75 ± 4.77 Oxt⁺ neurons/mouse project to the VTA; 37.50 ± 6.97 Oxt⁺ neurons/mouse project to the SNc; 2.25 ± 0.73 Oxt⁺ neurons/mouse project to both regions, *n* = 4 mice, one-way ANOVA with Tukey post hoc tests, *p* < 0.001) (Figures 1G and 1H).


For examining the projection of PVN Oxt neurons to midbrain DA regions, we relied on *Oxt*^{i-Cre} mice injected with a Cre-dependent recombinant adeno-associated viral vector (AAV) expressing eYFP into the PVN (Figure 1I). Here, 452/491 eYFP⁺ neurons were labeled for Oxt IF (3 mice) (Figure 1I). In addition to the well-described projections to amygdala and striatum (Knobloch et al., 2012), we observed eYFP⁺ axons in both the VTA and the SNc as well as in the lateral hypothalamus (Figure 1J; Figures S1C and S1D). We observed a mediolateral linear trend in the density of virally transduced axons, with the peak of expression in dorsolateral SNc and the poorly defined reticular area along its border (eYFP⁺ axonal density, VTA, 553.9 ± 86.4 mm/mm³; SNc, 783.7 ± 92.8 mm/mm³; reticular SNc, 1,515.9 ± 231.9 mm/mm³, *p* < 0.01, one-way ANOVA and post hoc test for linear trend, 3 mice) (Figures 1J and 1K; Figures S1C and S1D). Thus, both retrograde and anterograde tracing results confirm that PVN oxytocinergic efferents directly project to the VTA and SNc.

Distribution of Oxt-Sensing Receptors in the VTA and SNc

To evaluate which neurons in DA regions are capable of responding to Oxt, we investigated Oxt-sensing receptor expression

Figure 1. Targeting and Tracing Oxt Projections to the VTA and SNc

- (A) Left: confocal image of tdTomato (magenta) and immunofluorescent labeling (IF) for Oxt (green) in the PVN in *Oxt*^{i-Cre}; Ai14 mice. Merge shows signal overlap. Right: percentage of tdTomato⁺ neurons (magenta), Oxt⁺ by IF (green), and those co-localizing both signals (gray). *n* = 797 neurons, 3 mice.
- (B) Schematic of retrograde tracer experiments targeting VTA or SNc of *Oxt*^{i-Cre}; Ai14 mice.
- (C) Images of GRB injection sites in VTA/SNc of *Oxt*^{i-Cre}; Ai14 mice. TH, tyrosine hydroxylase.
- (D) Coronal brain section showing sparse GRB⁺ PVN cells after ipsilateral GRB_{VTA} targeting. Red, tdTomato; green, GRB; blue, nucleic stain.
- (E) Confocal images of the overlap between PVN tdTomato⁺ neurons (red) and GRB⁺ (green). Top: GRB_{VTA} injection; bottom: GRB_{SNc} injection. White solid circles mark double-labeled cells, and white dashed circles indicate tdTomato⁻/GRB⁺ cells.
- (F) Left: counts of GRB⁺ PVN neurons. Right: counts of GRB⁺/tdTomato⁺ cells. Circles, within mouse averages; *n* = 5 mice/group, Mann-Whitney test, *p* > 0.05.
- (G) Confocal images of the overlap between PVN Oxt neurons (blue), GRB⁺ cells after VTA targeting (green), and RRB⁺ cells after SNc targeting (red). Green circles mark Oxt⁺/GRB⁺ cells, red circles mark Oxt⁺/RRB⁺ cells, and yellow circles mark Oxt⁺/GRB⁺/RRB⁺ cells.
- (H) Average count of retro-labeled PVN Oxt⁺ neurons projecting to VTA and SNc. Gray, Oxt⁺/GRB⁺; purple, Oxt⁺/RRB⁺; yellow, triple-labeled neurons. One-way ANOVA with Tukey post hoc tests, ***p* < 0.001.
- (I) Left: AAV9-DIO-eYFP expression in the PVN of *Oxt*^{i-Cre} mice. Right: confocal image of eYFP⁺ neurons (green) and Oxt IF (magenta). Arrows point to double-labeled cells.
- (J) Left: oxytocinergic axons in midbrain DA regions. White dashed boxes mark locations of confocal images (right). Green, eYFP; magenta, TH IF.
- (K) Oxytocinergic axon coverage in the VTA and SNc. *n* = 3 mice, ***p* < 0.001 and **p* < 0.05, respectively, one-way ANOVA, post hoc test for linear trend. Error bars reflect SEM.



(legend on next page)

in the VTA and SNc. We analyzed tissue immunofluorescently double labeled for OxtR and the canonical marker of DA neurons, tyrosine hydroxylase (TH) (Figure 2A). An ~2-fold greater proportion of TH⁺ somata co-localized with OxtR⁺ puncta in the VTA versus in the SNc (~55% of VTA TH⁺ cells were OxtR IF⁺, but only ~26% in the SNc, $p < 0.05$, unpaired t test) (Figure 2B). Significantly fewer OxtR puncta were observed following IF in tissue where OxtR was conditionally knocked out from DAT⁺ neurons, confirming the specificity of receptor labeling. We also used a previously characterized OxtR antibody to confirm expression in the VTA and SNc (Marlin et al., 2015) (Figures S2A and S2B).

To refine and corroborate the results of IF assays, we carried out quantitative fluorescence in situ hybridization (FISH) assays for mRNA of *Oxtr* and *Avpr1a* in the VTA and SNc. DA and GABA neurons were identified by labeling for *Th* and *Slc32a1* (vesicular GABA transporter, *Vgat*) (Figures 2C and 2F). In the VTA, we observed co-localization of *Oxtr* with both *Th* and *Vgat* probes (proportion of *Th*⁺ cells expressing *Oxtr*, 6.86% ± 2.81%; proportion of *Vgat*⁺ cells expressing *Oxtr*, 8.38% ± 0.46%, $n = 3$ mice) (Figures 2D and 2E; Figure S2C). We observed a higher degree of co-localization of *Avpr1a* with *Th* and *Vgat* in the VTA (proportion of *Th*⁺ cells expressing *Avpr1a*, 25.70% ± 8.97%; proportion of *Vgat*⁺ cells expressing *Avpr1a*, 29.77% ± 6.02%) (Figures 2F–2H; Figure S2D). Since a subset of Oxt neurons co-releases Avp in lactating rats (Mezey and Kiss, 1991), we evaluated whether neurons in reporter mice expressing Cre recombinase under *Avp* promoter (*Avp*^{i-Cre}) are Oxt⁺. Despite the presence of *Avpr1a* in the VTA, we confirmed that <1% of PVN Oxt neurons (6/688 neurons, 2 mice) co-express Avp, consistent with a previous study (Knobloch et al., 2012) (Figures S2E and S2F).

In the SNc, few *Th*⁺ cells were positive for *Oxtr* or *Avpr1a* (*Oxtr*, 3.72% ± 0.39%; *Avpr1a*, 3.03% ± 1.14%); likewise, few *Vgat*⁺ cells co-localized with *Oxtr* or *Avpr1a* (*Oxtr*, 5.50% ± 0.16%; *Avpr1a*, 1.75% ± 0.53%) (Figures 2D–2H). Unexpectedly, we identified dense *Oxtr*⁺ puncta, distributed in the reticular part of the SNc, which co-localized with the *Vgat*⁺ cells (11.3% ± 2.04%) (Figures 2C–2E). These data support the notion that Oxt may regulate DA and GABA neurons in the VTA via OxtR and V1aR; furthermore, it may indirectly influence DA neurons in the SNc by modulating dorsolateral SNc reticular GABA neurons. This hypothesis is consistent with our observation that the greatest density of PVN oxytocinergic axons is in the most dorsolateral aspect of SNc (Figures 1J and 1K).

PVN Oxt Neurons Projecting to Midbrain DA Regions Are Parvocellular

Oxt neurons are classified into magnocellular and parvocellular subtypes, distinct in morphology, amount of Oxt production, and electrophysiological properties (Eliava et al., 2016; Luther and Tasker, 2000). Magnocellular Oxt neurons can be distinguished from parvocellular ones by a longer latency to spike in response to positive current injections and a transient outward rectification following relief from hyperpolarization (Eliava et al., 2016; Luther and Tasker, 2000). To determine the subtype that gives rise to these midbrain projections, we injected GRBs into either the VTA or the SNc of *Oxt*^{i-Cre}; Ai14 mice and recorded PVN Oxt neurons. PVN tdTomato⁺/GRB⁺ neurons in both GRB_{VTA}⁻ and GRB_{SNc}⁻ injected mice lacked this transient outward rectification and showed a short latency to spike. We found that they also had relatively small somata (Figures 3A and 3C). In contrast, *Oxt*⁺/GRB⁻ neurons with large somata showed transient outward rectification and had longer latencies to action potential firing when depolarized (Figures 3B and 3C). We further compared soma area of *Oxt*⁺/GRB⁺ and *Oxt*⁺/GRB⁻ neurons in confocal images from Figure 1 and found that *Oxt*⁺/GRB⁺ neurons have smaller somata (*Oxt*⁺/GRB⁺ neurons, 174.26 ± 6.58 μm²; *Oxt*⁺/GRB⁻ neurons, 215.12 ± 2.32 μm²; $p < 0.0001$, unpaired t test, $n = 79$ *Oxt*⁺/GRB⁺ neurons and 764 *Oxt*⁺/GRB⁻ neurons from 2 GRB_{VTA}⁻ and 2 GRB_{SNc}⁻ injected mice) (Figure 3D). Together with prior reports of differences between magnocellular and parvocellular neurons (Eliava et al., 2016; Luther and Tasker, 2000), our data suggest that PVN Oxt neurons projecting to VTA/SNc are primarily parvocellular.

Opposing Effects of Oxt on the Activity of VTA and SNc DA Neurons

To reveal the functional consequences of Oxt signaling for DA neurons, we recorded tdTomato⁺ neurons in slices prepared from *DAT*^{i-Cre}; Ai14 mice (Figure S3A). DA neurons tended to be tonically active, with firing rates (FRs) around 1–2 Hz and membrane potential “sag” in response to hyperpolarizing current injections (Guzman et al., 2009; Wanat et al., 2008) (Figures S3B and S3C). Consistent with the view that Oxt primarily acts at the G_{α_q} -coupled receptors in the central neural system (Grinevich et al., 2016), VTA DA neurons increased FR in the presence of 1 μM bath-applied Oxt, recovering back to baseline following removal of the peptide (control, 0.70 ± 0.16 Hz; Oxt, 0.94 ± 0.18 Hz; washout, 0.71 ± 0.18 Hz; $p < 0.05$, one-way ANOVA with Tukey post hoc tests, $n = 22$ neurons, 15 mice).

Figure 2. Oxt-Sensing Receptor Protein and Transcript Distribution in the VTA and SNc

- (A) Confocal images of TH (white) and OxtR IF (cyan) in VTA and SNc.
 (B) Quantification of OxtR⁺ puncta co-localization with TH⁺ neurons. $n = 780$ VTA TH⁺ neurons, 564 SNc TH⁺ neurons. Inset: proportion of neurons co-localized with OxtR⁺ puncta. $n = 6$ sections, 2 mice; Unpaired t test, $p < 0.05$. Error bars reflect SEM.
 (C) Fluorescence in situ hybridization (FISH) images in the VTA (top) and dorsolateral SNc reticular area (SNc-r, bottom). *Th* (magenta), vesicular GABA transporter (*Slc32a1/Vgat*, orange), and *Oxtr* (cyan).
 (D) Quantitative analysis of *Th*, *Vgat*, and *Oxtr* co-expression by FISH in the VTA (left), SNc (middle), and SNc-r area (right). The number of puncta per cell for either *Th* (pink triangles) or *Vgat* (orange squares) against *Oxtr* (both). Magenta and orange lines mark the cutoffs used to classify a cell as *Th*⁺ or *Vgat*⁺; blue line, the cutoff for *Oxtr* ($n = 3$ mice).
 (E) Cumulative frequency distribution of *Oxtr*⁺ neurons in regions of interest (ROIs) analyzed for *Th*⁺ cells (top) and *Vgat*⁺ cells (bottom).
 (F) Same as (C), but for *Avpr1a*.
 (G) Quantitative analysis of *Th*, *Vgat*, and *Avpr1a* co-expression by FISH in VTA (left) and SNc (right).
 (H) Cumulative frequency distributions of *Avpr1a*⁺ neurons in ROIs analyzed for *Th*⁺ cells (left) and *Vgat*⁺ cells (right).




Figure 3. Parvocellular Oxt Neurons of the PVN Project to Midbrain DA Regions

(A) Soma morphology and physiological response properties of an SNc-projecting PVN Oxt neuron filled with Alexa 488 and imaged on a two-photon laser-scanning microscope.

(B) Same as (A), but for a GRB⁻ Oxt neuron.

(C) Left: latency to first spike in response to a 40 pA current injection. Right: soma size. Green/black circles mark neurons shown in (A) and (B). $n = 6$ neurons for Oxt⁺/GRB⁺ group, 8 neurons for Oxt⁺/GRB⁻ group, 5 mice (2 and 3 for RB VTA and SNc targeting), unpaired t test, * $p < 0.05$ and ** $p < 0.01$.

(D) Summary of PVN Oxt neuron soma size from retrograde tracing experiments. $n = 79$ neurons for Oxt⁺/GRB⁺ group and 764 neurons for Oxt⁺/GRB⁻ group from 4 mice (2 VTA targeted and 2 SNc targeted), unpaired t test, *** $p < 0.001$. Error bars reflect SEM.

(Figures 4A and 4C). Oxt-responsive neurons were defined here and throughout as those exhibiting at least a 20% change from baseline activity level in response to Oxt application—for these neurons, Oxt-mediated enhancement in firing was approximately 2-fold ($210.5\% \pm 22.0\%$ of baseline activity) (Figure 4C). The observed effect on tonic activity in the presence of Oxt was accompanied by a modest decrement in afterhyperpolarization and action potential height (Figure S3D). To confirm that this modulation did not depend on the intracellular recording configuration, we recapitulated this observation using cell-attached recordings of DAT⁺ neurons (Figure S3E).

In contrast, the FRs of SNc DA neurons during Oxt application showed a more heterogeneous set of responses, correlated with the mediolateral positions of the neurons. Medial SNc DA neurons did not show a significant FR modulation in response to Oxt application (control, 0.87 ± 0.29 Hz; Oxt, 1.02 ± 0.27 Hz; washout, 0.80 ± 0.30 Hz; one-way ANOVA, $p = 0.270$, $n = 8$ neurons, 8 mice) (Figure 4D). However, lateral SNc DA neurons

reversibly decreased FR in the presence of Oxt (control, 0.91 ± 0.18 Hz; Oxt, 0.49 ± 0.14 Hz; washout, 0.91 ± 0.22 Hz; $p < 0.05$, one-way ANOVA with Tukey post hoc tests, $n = 14$ neurons, 12 mice) (Figures 4B and 4D). In the lateral SNc, Oxt decreased the FR of Oxt-sensitive DAT⁺ neurons by ~70% ($31.8\% \pm 7.9\%$ of baseline firing rate) (Figure 4D). For lateral SNc DA neurons recorded in cell-attached mode, spontaneous activity was also inhibited by Oxt (Figure S3F). Oxt narrowed the width and increased the height of SNc DA neuron action potentials (Figure S3D). Regardless of the anatomical distinction of VTA versus SNc, the lateral location of a DA neuron predicted a decrement in FR in response to Oxt, while a medial location was associated with an increase in FR (Figure 4E). This divergent neuromodulatory regulation of tonic activity was largely sex invariant (VTA FR change: ♂ 0.23 ± 0.10 Hz, ♀ 0.24 ± 0.09 Hz, $p = 0.919$, unpaired t test, $n = 11$ neurons/group; SNc FR change: ♂ -0.71 ± 0.23 Hz, ♀ -0.25 ± 0.12 Hz, $p = 0.101$, unpaired t test, $n = 12$ and 10 neurons/group).




Figure 4. VTA and SNc DA Neuron Activity Is Differentially Modulated by Oxt

(A) Top: FR histogram (10 s bins) from a spontaneously active VTA DAT⁺ neuron with bath application of 1 μ M Oxt (blue bar). Bottom: traces correspond to time points 1–3 above.

(B) Same as (A), but in the SNc.


(C) Left: Oxt-induced FR increase in VTA DA neurons. Green circles, Oxt-responsive neurons that increased activity. The larger circle marks neuron shown in (A). Paired t test on full dataset, *** $p < 0.001$, $n = 22$ neurons, 15 mice. Right: normalized effect magnitudes for Oxt-responsive neurons.

(D) Left: Oxt-induced FR decrease in lateral SNc DA neurons. Yellow circles, Oxt-responsive neurons that decreased activity. The larger yellow circle marks neuron shown in (B). Paired t test, ** $p < 0.01$, $n = 14$ neurons, 12 mice. Middle: normalized effect magnitudes for Oxt-responsive lateral SNc neurons that decreased activity with Oxt. Right: medial SNc DAT⁺ neuronal population showed a mixed response to Oxt. Paired t test, $p = 0.332$, $n = 8$ neurons, 8 mice.

(E) Spatial distribution of spontaneously active DA neurons, color coded for significant FR responses to Oxt (defined in text). Green reflects increased FR; yellow, decreased FR; black, unchanged. Right, middle: collapsing across rostro-caudal position reveals a mediolateral gradient in Oxt responsiveness. Larger green and yellow circles mark the location of neurons shown in (A) and (B). Right, bottom: Oxt-induced FR changes correlate with the mediolateral position of DA neurons (grouped by recording location). Red line, second order polynomial fit.

(F) Left: traces from a VTA DA neuron during baseline, WAY267464 application, and washout. Middle: FR before and in WAY267464. Coloring as in (E). Larger circle marks the example neuron. Paired t test, * $p < 0.05$, $n = 12$ neurons, 8 mice. Right: normalized effect magnitudes for Oxt-responsive neurons that enhanced FR in WAY267464.

(G) Same as (F), but in the SNc. Paired t test, * $p < 0.05$, $n = 13$ neurons, 8 mice. Error bars reflect SEM.



(legend on next page)

To validate the results of Oxt application experiments, we used a specific and selective OxtR agonist WAY267464 while recording VTA or lateral SNc DAT⁺ neurons. Just as for Oxt, bath-applied WAY267464 increased the FR of VTA DA neurons (control, 1.39 ± 0.19 Hz; WAY267464, 1.61 ± 0.23 Hz; $p < 0.05$, paired t test, $n = 12$ neurons, 8 mice) and decreased the FR of lateral SNc DA neurons (control, 1.52 ± 0.19 Hz; WAY267464, 1.26 ± 0.28 Hz; $p < 0.05$, paired t test, $n = 13$ neurons, 8 mice) (Figures 4F and 4G). These FR changes corresponded to a $153.7\% \pm 25.7\%$ enhancement for agonist-responsive DAT⁺ neurons in the VTA and an over 2-fold decrease in activity for SNc DAT⁺ neurons ($38.7\% \pm 4.5\%$ of baseline FR).

Optogenetic Stimulation of Oxt Terminals Differentially Regulates VTA and SNc DA Neurons

Next, we investigated whether endogenous Oxt release is sufficient to regulate the activity of midbrain DA neurons. To selectively control release, we relied on Oxt^{i-Cre} mice injected with a Cre-dependent AAV expressing a fusion of ChR2 and either mCherry (ChR2-mCherry) or eYFP (ChR2-eYFP) into the PVN (Figure 5A). A single medial injection provided dense and specific coverage of PVN Oxt neurons: 712/839 ChR2⁺ neurons from 5 mice overlapped with Oxt IF (Figure 5A; Figures S4A and S4B). Virally labeled axons were observed in the VTA and SNc, corroborating anterograde mapping experiments. PVN ChR2⁺ neurons reliably responded to pulses of 470 nm light (Figure 5B). Since high-frequency stimulation is required to induce Oxt release (Knobloch et al., 2012), to evoke axonal release of Oxt, we used trains of 10-ms-long light pulses at 50 Hz delivered in 15-s-long trains.

In this configuration, Cre recombinase expression is reserved for activating Oxt fibers and is not available for identifying DA neurons. To reliably target DA cells without fluorescence reporter expression, we first recorded 45 DAT⁺ neurons in the VTA and SNc of DAT^{i-Cre}; Ai14 mice to set electrophysiological and morphological criteria for defining a DA neuron. Then, we evaluated these criteria in unlabeled VTA and SNc neurons in Oxt^{i-Cre} mice, identified as DA⁺ based on their larger soma size, the presence of a sag potential, and their low tonic firing (Figures S3A–S3C), and validated with post hoc TH immunolabeling

(Figure S4C). Others also report that DA neurons in mice can be identified based on their response properties and morphology (Ungless and Grace, 2012; Wanat et al., 2008).

Consistent with the results of bath application experiments, we observed an enhancement of FR in VTA DA neurons when optogenetically activating oxytocinergic axons (baseline, 1.19 ± 0.18 Hz; light, 1.37 ± 0.19 Hz; after, 1.13 ± 0.19 Hz; $p < 0.01$, one-way ANOVA with Tukey post hoc tests, $n = 16$ neurons, 12 mice) (Figure 5C), in contrast to a decrement in FR in lateral SNc DA neurons during light stimulation (baseline, 1.21 ± 0.16 Hz; light, 0.90 ± 0.19 Hz; after, 1.04 ± 0.17 Hz; $p < 0.01$, one-way ANOVA with Tukey post hoc tests, $n = 16$ neurons, 10 mice) (Figure 5D). For light-sensitive VTA DA neurons, defined above (20% FR modulation), stimulation increased FR to $132.8\% \pm 4.7\%$ of baseline (Figure 5C); in the SNc, activity in responsive neurons decreased to $37.1\% \pm 7.6\%$ of baseline (Figure 5D). For DA neurons lacking spontaneous firing, light stimulation elicited analogous opposing changes in resting membrane potential (Figures S4D and S4E). These Oxt-induced voltage changes are physiologically relevant, as small differences in membrane potential translate into relatively large FR changes for DA neurons (Figure S4F).

DA neurons provide the primary output of the VTA and SNc, respectively targeting the vStr to regulate reward and the dStr to modulate locomotion (Howe and Dombeck, 2016; Watabe-Uchida et al., 2012), but they also innervate multiple other regions (Beier et al., 2015; Walsh and Han, 2014). To determine which populations of DA neurons are modulated by Oxt, we combined GRB injections and viral transduction of ChR2. GRBs were injected into dStr or vStr of Oxt^{i-Cre} mice transduced with AAV-ChR2-mCherry in the PVN, and we recorded GRB⁺ neurons from the VTA and SNc, with light stimulation as above (Figures 5E and 5F). Consistent with previous studies (Lammel et al., 2011; Watabe-Uchida et al., 2012), most GRB⁺ neurons observed in midbrain DA regions were located in the VTA following vStr injections but in the SNc after dStr GRB targeting (Figure S4G). In response to light activation of Oxt fibers, GRB⁺ VTA DA neuron excitability increased, based on measurements of FR or membrane potential (6/11 neurons, 4 mice) (Figure 5E). On the other hand, most GRB⁺ SNc DA neurons showed

Figure 5. Optogenetic Activation of Oxytocinergic Fibers Leads to Opposite Activity Changes in VTA and SNc DA Neurons

(A) Left: schematic of viral transduction strategy to express ChR2 in PVN Oxt neurons of Oxt^{i-Cre} mice. Middle: ChR2-mCherry (red) expression confirmed with Oxt IF (white). Right: confocal images of axonal mCherry signal in VTA (top) and SNc (bottom), in proximity of DA neurons (TH IF, blue).

(B) Current-clamp trace of an mCherry⁺ PVN neuron in response to 5-ms-long 470 nm light pulses (10 Hz).

(C) Left: current-clamp trace from a VTA DA neuron (DA_n). Light stimulation of ChR2-expressing Oxt terminals in the VTA with 10-ms-long light pulses delivered at 50 Hz (15 s) reversibly increased FR. Middle: VTA DA neuron FR changes. Paired t test on full dataset, *** $p < 0.001$, $n = 16$ neurons, 12 mice. Green, DA neurons that enhanced their FR. Larger circle marks the neuron shown in left panel. Right: normalized effect magnitudes for light-responsive neurons.

(D) Same as (C), but in the SNc. Yellow, light-sensitive neurons that decreased FR; larger circle, neuron shown in left panel. Paired t test, ** $p < 0.01$, $n = 16$ neurons, 10 mice.

(E) Left: schematic depicting the double injection of GRB into striatum and an AAV to express ChR2 in the PVN of Oxt^{i-Cre} mice. Right: the majority of VTA DA neurons projecting to vStr increased activity in response to light, based on FR or membrane potential changes. $n = 11$ neurons, 4 mice.

(F) Same as (E), but for the SNc and dStr. Right: the majority of SNc DA neurons projecting to dStr decreased activity in response to light. $n = 13$ neurons, 3 mice.

(G) Top: schematic of the experimental design. Left: VTA DA neuron FR changes in response to light stimulation before and during the application of OxtR antagonist (L368,899). $n = 6$ neurons, 6 mice. Right: same as in left panel, but for the SNc. $n = 9$ neurons, 6 mice. * $p < 0.05$, paired t test, ## $p < 0.01$, t test versus baseline.

(H) Left: FR changes in all VTA DA neurons in standard ACSF ($n = 16$ neurons, 12 mice) and in the presence of blockers of fast synaptic transmission (CPP, NBQX, and SR95531, $n = 8$ neurons, 6 mice). Right: as in left panel, but for the SNc (ACSF, $n = 16$ neurons, 10 mice; with blockers, $n = 23$ neurons, 17 mice). n.s., $p > 0.05$, unpaired t test; ## $p < 0.01$, t test versus baseline. Error bars reflect SEM.

reversible inhibition of activity (10/13 neurons, 3 mice) (Figure 5F). These results indicate that PVN Oxt projections are poised to shift the overall balance of DA tone from SNc to VTA-specific sources in downstream targets of DA neurons.

Oxt Directly Enhances the Activity of VTA DA Neurons

Optogenetic control of Oxt release enabled selection of DA neurons displaying specific modulatory responses for further dissection of the underlying mechanisms. In the VTA, light-evoked FR increases in DA neurons were abolished by the OxtR antagonist L368,899, used at concentrations that should block both OxtR and, to a lesser degree, V1aR (FR change_{ACSF}, 0.26 ± 0.05 Hz; FR change_{L368,899}, 0.01 ± 0.06 Hz, $p < 0.05$, paired t test, $n = 6$ neurons, 6 mice) (Figure 5G). Light-evoked changes in activity persisted in the presence of a cocktail of antagonists for glutamatergic and GABAergic receptors, supporting a direct, monosynaptic effect of Oxt release on VTA DA neurons (Figure 5H).

Oxt Indirectly Inhibits SNc DA Neurons via Local GABA Neurons

Several lines of evidence suggest that in the SNc, rather than directly activating Oxt-sensing receptors on DA neurons, Oxt acts through distinct neural circuits or receptor mechanisms. First, we found few OxtRs in SNc DA neurons (Figure 2). Second, the observed inhibition of SNc DA neuron activity by Oxt is inconsistent with canonical $G\alpha_q$ -mediated effects found in other brain areas (Huber et al., 2005; Zheng et al., 2014). Finally, light-evoked inhibition of SNc DA neurons was longer lasting relative to the opposite valence modulation observed in the VTA (Figure S5C). Consistent with the possibility of multi-receptor mechanisms, OxtR antagonist alone was insufficient to block light-evoked inhibition of SNc DA neurons (FR change_{ACSF} -0.27 ± 0.04 Hz; FR change_{L368,899} -0.40 ± 0.08 Hz, $p = 0.166$, paired t test, $n = 9$ neurons, 6 mice) (Figure 5G).

If Oxt-induced inhibition of SNc DA neurons is mediated by local circuits, it may require fast neurotransmission. To test this hypothesis, we carried out additional pharmacological dissection of light-evoked activity changes in stimulation-responsive DA neurons. Blocking fast neurotransmission with CPP, NBQX, and SR95531 did not abolish light-mediated inhibition of SNc DA neuron activity (Figure 5H). However, in the absence of fast transmission, the inhibition became more transient, similar in timescale to the enhancement of tonic activity in the VTA (Figures S5A–S5C). With fast transmission blocked, light-evoked inhibition persisted in the presence of additional selective blockers for each of the following receptors individually: Oxt (L368,899, -0.33 ± 0.09 Hz, $p < 0.01$, t test versus baseline, $n = 7$ neurons, 5 mice), V1a (SR49059, -0.32 ± 0.06 Hz, $p < 0.01$, $n = 6$ neurons, 4 mice), and Drd2 (L741,626, -0.30 ± 0.08 Hz, $p < 0.01$, $n = 9$ neurons, 7 mice) (Figure 6A). However, GABA(B)R antagonist (CGP54626) abolished light-driven inhibition of activity in SNc DA neurons (FR change_{control}, -0.31 ± 0.05 Hz; FR change_{CGP54626}, -0.08 ± 0.05 Hz, $p < 0.01$, paired t test; $n = 10$ neurons, 9 mice). FR change_{CGP54626} did not differ from baseline ($p = 0.122$, t test versus baseline) (Figures 6A and 6B), raising the likelihood of a disynaptic effect via GABA neurons. To evaluate the role of GABAergic neurons in mediating oxytocinergic

regulation of SNc DA neurons, we repeated these experiments with synaptic signaling intact. Light-evoked hyperpolarization of SNc DA neurons was diminished in the presence of GABA(A)R and GABA(B)R antagonists (membrane potential change_{control}, -2.46 ± 0.37 mV; with blockers, -0.53 ± 0.22 mV; $p < 0.01$, paired t test, $n = 9$ neurons, 5 mice) (Figure 6C).

We next sought to reveal the basis for oxytocinergic inhibition of SNc DA neuron activity using pharmacological blockers for GPCRs and signaling cascades. Since multiple Oxt-sensing receptors are expressed in the SNc (Skuse and Gallagher, 2009), we recorded SNc DA neurons in the presence of OxtR and Avpr (V1aR and V1bR) antagonists. Each blocker alone had no significant effect on Oxt-mediated inhibition of SNc DA neuronal activity (Figures 6A and 6D). Yet, together, these antagonists sufficed to fully block the modulatory effect of Oxt (control membrane potential change, -2.67 ± 0.45 mV; with blockers, 0.05 ± 0.28 mV; $p < 0.01$, paired t test, $n = 5$ neurons, 3 mice) (Figure 6E). Blocking $G_{\beta\gamma}$ subunit-dependent signaling with gallein also prevented Oxt-driven effects, confirming a GPCR-based mechanism of SNc inhibition by Oxt (membrane potential change_{control} -2.49 ± 0.41 mV; blockers, -0.45 ± 0.26 mV; $p < 0.01$, paired t test, $n = 6$ neurons, 4 mice) (Figure 6F).

Our previous experiments do not distinguish whether GABA-mediated inhibition stems from local interneurons or directly from Oxt projections. Although we did not observe light-evoked inhibitory currents (IPSCs) in DA neurons (data not shown), we evaluated the possibility of GABA co-release from Oxt neurons. We measured the overlap between Oxt IF and tdTomato⁺ neurons in the PVN of *Vgat*^{i-Cre}; tdTomato mice and found 0/341 double-labeled cells in 2 mice (Figure 7A). Therefore, Oxt neurons are unlikely to co-release GABA. Next, we confirmed the basis for disynaptic oxytocinergic modulation via local inhibitory interneurons by recording spontaneous IPSCs (sIPSCs) in midbrain DA regions. For VTA DA neurons, light activation of Oxt fibers enhanced sIPSC frequency (baseline, 11.09 ± 2.56 Hz; light, 16.59 ± 4.87 Hz; $p < 0.05$, paired t test, $n = 6$ neurons, 2 mice) but did not change their amplitude (baseline, 41.91 ± 5.25 pA; light, 40.66 ± 5.78 pA; $p = 0.691$) (Figures 7B and 7C). This disynaptic inhibition was insufficient to override the direct, activity-enhancing modulation of VTA DA neurons by Oxt (Figure S5D). For SNc DA neurons, light activation of Oxt fibers also significantly increased sIPSC frequency and shortened inter-event intervals (baseline, 18.92 ± 3.91 Hz; light, 21.66 ± 4.46 Hz; $p < 0.05$, paired t test, $n = 6$ neurons, 2 mice) without altering sIPSC amplitude (baseline, 32.55 ± 5.47 pA; light, 31.75 ± 5.55 pA; $p = 0.131$) (Figures 7D and 7E). Direct bath application of Oxt also increased sIPSC frequency and shortened the inter-event intervals in both VTA and SNc DA neurons (VTA: baseline, 8.75 ± 2.28 Hz; Oxt, 10.11 ± 2.53 Hz; $p < 0.05$, paired t test, $n = 5$ neurons, 4 mice; SNc: baseline, 12.78 ± 7.94 Hz; Oxt, 13.45 ± 7.99 Hz; $p < 0.05$, paired t test, $n = 5$ neurons, 4 mice) (Figures 7F and 7G).

Analogously, TH⁺ neurons in the VTA and SNc, presumed to be GABA interneurons, showed an enhancement of FR in response to light stimulation (VTA: FR_{control} 5.45 ± 0.61 Hz, light, 6.69 ± 0.73 Hz, recovery, 5.85 ± 0.91 Hz; $p < 0.05$, one-way ANOVA with Tukey post hoc tests, $n = 7$ neurons, 4 mice; SNc: FR_{control} 3.24 ± 0.59 Hz, light, 3.78 ± 0.69 Hz, recovery, 3.05 ± 0.65 Hz;




Figure 6. Optogenetic Stimulation of Oxt Terminals Inhibits SNc DA Neurons through Diverse Circuit and Receptor Mechanisms

(A) Top: schematic of the experimental design. Bottom: lateral SNc DA neuron FR changes in response to light stimulation in the presence of blockers of fast synaptic transmission and antagonists for the following receptors: OxtR (L368,899, $n = 7$ neurons, 5 mice), V1aR (SR49059, $n = 6$ neurons, 4 mice), Drd2 (L741,626, $n = 9$ neurons, 7 mice), and GABA(B)R (CGP54626, $n = 10$ neurons, 9 mice). t test versus baseline, $**p < 0.01$.

(B) Traces from one SNc DA neuron before and during the application of GABA(B)R antagonist CGP54626 combined with light stimulation.

(C) In ACSF, light-mediated hyperpolarization of SNc DA neurons is blocked by GABA(A)R and GABA(B)R antagonists. Gray line, raw data; black line, smoothed. Paired t test, $**p < 0.01$, $n = 9$ neurons, 5 mice.


(D) Same as (C), but for the V1bR antagonist. Paired t test, $p = 0.692$, $n = 6$ neurons, 3 mice.

(E) Light-induced hyperpolarization is fully blocked by a combination of OxtR, V1aR, and V1bR antagonists (L368,899, SR49059, and SSR149415). Gray line, raw data; black line, smoothed. Paired t test, $**p < 0.01$, $n = 5$ neurons, 4 mice.

(F) Same as (E), but for antagonist of G protein $\beta\gamma$ subunit-dependent signaling, gallein. Paired t test, $**p < 0.01$, $n = 7$ neurons, 4 mice. Error bars reflect SEM.

$p < 0.05$, one-way ANOVA with Tukey post hoc tests, $n = 12$ neurons, 7 mice) (Figures 7H and 7I). Light-induced FR changes were attenuated by OxtR antagonist (FR change_{control}, 1.23 ± 0.34 Hz;

with blocker, 0.42 ± 0.38 Hz; $p < 0.05$, paired t test, $n = 5$ neurons, 5 mice) (Figure 7J). Although Oxt has the capacity to act via $G_{i/o}$ proteins to inhibit neuronal activity (Busnelli et al., 2012; Eliava



(legend on next page)

et al., 2016; Stoop, 2012), our results support a model where Oxt activates multiple GPCRs to indirectly inhibit SNc DA neurons via GABA neurons. A similar dampening in the VTA is compensated by direct Oxt action on DA neurons, yielding an overall activity enhancement.

DISCUSSION

Behavioral and anatomical studies have suggested the possibility of oxytocinergic modulation of midbrain DA neurons, but knowledge of the circuits and the physiological consequences of this potential modulation have remained elusive. Here, we anatomically and functionally characterize hypothalamic oxytocinergic projections that regulate midbrain DA neuron activity, supporting a direct interaction between neurohypophyseal systems and DA circuits (Figure 8). This oxytocin source derives from parvocellular neurons of the PVN. In the VTA, Oxt directly increases the FR of most DA neurons. In the SNc, by contrast, oxytocin hyperpolarizes membrane potential, causing a net decrease in FR. The opposition of these net modulatory effects stems from differences in receptor expression and circuit connectivity. Oxytocin-sensing receptors are expressed on both DA and GABA neurons in the VTA. However, few oxytocin-sensing receptors are expressed on SNc DA neurons, unlike the population of nearby GABA cells.

This biased modulation, mediated by oxytocin-sensing receptor gradients and differences in circuit organization, demonstrates a mediolateral gradient of responsiveness to Oxt in midbrain DA regions. Other mediolateral cellular gradients are well established in this region. Lateral DA regions develop earlier than medial areas (Bayer et al., 1995), and they degenerate more quickly and intensely than VTA DA neurons during the clinical progression of Parkinson's disease (PD) (Duke et al., 2007). Since PVN Oxt neurons are also vulnerable in PD (Purba et al., 1994), future studies are needed to explore the possibility that a loss of oxytocinergic dampening in the SNc contributes to DA neuron damage and the pathophysiology of PD.

Differences in expression patterns of receptors provide a simple mechanistic framework for mediating biased responses. The single OxtR shares high levels of sequence alignment with the three AvpRs (Mouillac et al., 1995). Compared to the

activity of Oxt via OxtR, its effect on V1aR and V2R is about 1% of full activity (Manning et al., 2008; Tribollet et al., 1988). Still, Oxt action at V1aR can mediate peripheral analgesic effects (Qiu et al., 2014; Schorscher-Petcu et al., 2010). In addition, the activity of Oxt at V1bR is involved in the secretion of insulin and glucagon from the pancreatic β cells and adrenocorticotrophic hormone from the anterior pituitary (Fujiwara et al., 2007; Lee et al., 1995). Altogether, previous studies have shown that Oxt can act at AvpRs, with physiological consequences for the organism. However, it had remained unclear whether Oxt release exerts physiologically meaningful effects through central AvpRs. We relied on optogenetic stimulation of Avp-negative Oxt terminals and found that a small subset of light-evoked modulatory effects require AvpR antagonists for full blockade. So, despite the distinctions of nomenclature, the actions of Oxt and Avp are likely intertwined. In addition to Oxt acting on AvpRs in midbrain DA regions, Avp could also directly modulate these areas, since Avp neurons project to the VTA (Beier et al., 2015). Our retrograde labeling experiments indicate that only ~20% of VTA/SNc-targeting PVN neurons are Oxt⁺. Future studies are necessary to reveal the full scope of PVN peptides that could alter DA release and to identify local interactions of these molecules.

The biased oxytocinergic signaling that we observed is reminiscent of other subcortical systems modulated by neurohypophyseal peptides. The spatial separation of nearby neurons expressing OxtR and V1aR appears in other brain regions (Stoop, 2012; Veinante and Freund-Mercier, 1997). For example, oxytocin and vasopressin projections match local, segregated patterns of OxtR and V1aR expression in the central amygdala (Huber et al., 2005). However, our data suggest the sufficiency of Oxt release for physiological effects via multiple GPCRs, distributed in area-specific and cell-class-specific ways. This distribution of neurohormone receptors in the VTA/SNc sets up midbrain DA regions to be controlled by either Oxt or Avp or both. The close spatial proximity between oxytocin-sensing receptors in midbrain DA regions and Oxt⁺ axons of PVN neurons supports the likelihood of direct synaptic effects, as in the central amygdala (Knobloch et al., 2012). Yet, Oxt neurons are capable of dendritic release of oxytocin (Landgraf and Neumann, 2004; Ludwig et al., 2002) and can signal through microvolume diffusion (Fuxe et al., 2012), which could also contribute to the regulation of DA neuron activity in vivo.

Figure 7. Optogenetic Activation of Oxytocinergic Axons Enhances the FR of VTA and SNc GABA Neurons

- (A) Confocal image of *Vgat*-tdTomato reporter expression in the PVN (magenta) and Oxt IF (green). No signal overlap was observed ($n = 341$ neurons, 2 mice).
 (B) Traces from a VTA DA neuron showing spontaneous IPSCs (sIPSCs) before (top) and during (bottom) light stimulation (in CPP and NBQX).
 (C) Light-evoked changes in sIPSC inter-event intervals (IEIs) and frequency (left), with no change in sIPSC amplitude (right). Insets show normalized sIPSC frequency and amplitude. Paired t test, $*p < 0.05$, $n = 6$ neurons, 2 mice.
 (D) Same as (B), but in the SNc.
 (E) Same as (C), but in the SNc.
 (F) Traces from a VTA DA neuron showing sIPSCs before (top) and during (bottom) bath application of Oxt (in CPP and NBQX).
 (G) Oxt decreases sIPSC IEI and increases sIPSC frequency in VTA (left) and SNc (right) DA neurons. Insets show normalized sIPSC frequency. Paired t test, $*p < 0.05$, $n = 5$ neurons, 4 mice/group.
 (H) Top: a TH⁺ neuron in the SNc, filled with Alexa 594 (left), fixed and labeled with TH IF (middle). Bottom: traces from an SNc TH⁺ neuron in response to light stimulation of Oxt fibers.
 (I) FR changes in VTA (left) and SNc (right) TH⁺ neurons in response to light stimulation. One-way ANOVA (Tukey post hoc), $*p < 0.05$, $n = 7$ VTA neurons, 4 mice, $n = 12$ SNc neurons, 7 mice.
 (J) In SNc TH⁺ neurons, light stimulation-induced FR increase is diminished by OxtR antagonist L368,899. Paired t test, $*p < 0.05$, $n = 5$ neurons, 5 mice.




Figure 8. Model of Oxytocinergic Regulation of VTA and SNc Neurons

Parvocellular Oxt neurons located in the PVN, but not the SON, directly project to the VTA and SNc. In the VTA, Oxt-sensing receptors are expressed on DA and GABA neurons. The net modulatory effect of Oxt release is an increase in VTA DA neuron activity. In the SNc, few Oxt-sensing receptors are expressed on DA neurons, while a population of nearby GABA cells strongly express them. The consequence of these differences is a net decrease of SNc DA neuron activity in response to Oxt.

extensive oxytocin circuitry that biases the activity of VTA and SNc DA neurons in a region-specific and opposing manner. The present work represents the first observation that direct, evoked oxytocin release from Oxt⁺ axons modulates the activity of VTA and SNc DA neurons.

Parvocellular Oxt neurons of the PVN give rise to substantial central projections, including those targeting limbic and brainstem regions (Eliava et al., 2016; Insel and Shapiro, 1992). The Oxt neurons we observe projecting to the VTA and SNc comprise largely separate groups. It remains to be determined whether they send additional projections outside the DA midbrain. Future studies are also needed to reveal the potential interactions between sex and Oxt control of reward circuitry, as neurohypophyseal systems have sex-specific functions (Crowley, 2015; Young and Wang, 2004), and magnocellular Oxt neurons prime labor induction and lactation (Gimpl and Fahrenholz, 2001; Insel and Shapiro, 1992). Our anatomical and physiological data on oxytocinergic regulation of midbrain DA systems support relative sex invariance in structure and function. Thus, this neuromodulatory input may control ethologically significant behaviors shared by males and females. When could these behaviors manifest? In mice, OxtR expression changes over the course of the first several weeks of life (Mitre et al., 2016). We found that oxytocinergic regulation of DA neurons is consistent across several weeks of postnatal development (~P21–P50). Future studies will determine when the critical functions of maternal circulating oxytocin (Mazzuca et al., 2011; Tyzio et al., 2006) are replaced by endogenously produced peptide.

In addition to delineating the circuits for direct oxytocinergic control over midbrain DA systems, our study highlights the likely importance of local GABA neurons near the SNc for the regulation of SNc DA neuron activity. The main, well-established GABAergic projection to the SNc originates in the substantia nigra pars reticulata (SNr). However, SNr neurons do not contain significant amounts of oxytocin-sensing receptors or Oxt axons. Instead, we found a high concentration of *Oxtr* transcripts just dorsolateral to the SNc. While this reticular region is poorly defined anatomically, the GABAergic population that resides there may interact with DA neurons of the SNc (Hebb and Robertson, 2000). It has been previously shown that oxytocin application can directly activate a subset of neurons, including those in the medial VTA via $G\alpha_q$ -coupled OxtRs (Tang et al., 2014), but our study details

What is the function of oxytocinergic regulation of midbrain DA systems? Given the distributed projections of DA neurons, one compelling hypothesis is that oxytocin and DA signals cooperate to refine salient environmental cues. Indeed, studies suggest that oxytocin acts in the auditory cortex and olfactory bulb to improve the processing of salient sensory input (Marlin et al., 2015; Oettl et al., 2016). DA input from the VTA to the auditory cortex serves an analogous function (Bao et al., 2001; Lou et al., 2014). We find that optogenetic activation of midbrain Oxt fibers biases the activity of VTA and SNc DA neurons that send canonical ventral and dorsal projections. This modulation could regulate behavior by altering tonic activity of DA neurons and biasing striatal DA tone. Based on the known effects of striatal oxytocin on social interactions and pair-bonding behavior in the prairie vole and other rodents (Burkett et al., 2016; Dölen et al., 2013; Gao et al., 2016), one intriguing possibility is that oxytocin biases DA tone to enhance socially rewarding interactions at the expense of exploratory locomotion. Consistently, oxytocin microinjections into the VTA enhance the preference for social interactions (Song et al., 2016) and decrease locomotion when delivered into the SNc (Angioni et al., 2016). Our study delineates the circuit and receptor logic that supports direct oxytocin-mediated control of DA circuitry, adding a new motif to the complex picture of interactions between the basal ganglia and the neurohypophyseal systems that powerfully regulate social behaviors critical for species survival.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE
- CONTACT FOR REAGENT AND RESOURCE SHARING
- EXPERIMENTAL MODEL AND SUBJECT DETAILS
 - Mouse Strains and Genotyping

● METHOD DETAILS

- Stereotaxic Intracranial Injections and Fiber Optic Implants
- Tissue Processing, Immunohistochemistry, and Imaging
- Quantitative Fluorescence In Situ Hybridization (FISH)
- Image Analysis
- Acute Slice Preparation and Electrophysiology
- 2-Photon Laser Scanning Imaging
- Pharmacology

● QUANTIFICATION AND STATISTICAL ANALYSIS

SUPPLEMENTAL INFORMATION

Supplemental Information includes five figures and can be found with this article online at <http://dx.doi.org/10.1016/j.neuron.2017.06.003>.

AUTHOR CONTRIBUTIONS

L.X. and Y.K. designed the experiments. L.X. performed *ex vivo* electrophysiology. L.X. and J.N. carried out stereotaxic surgery. M.F.P. performed FISH experiments and analysis. L.X., J.N., and T.L. carried out immunohistochemistry experiments and analysis. L.X. and Y.K. wrote the paper, with extensive contributions from all authors.

ACKNOWLEDGMENTS

We thank Lindsey Butler for genotyping and colony management, Kozorovitskiy lab members for comments, and Dr. Robert Froemke for the OxtR-2 antibody. We thank the Northwestern University Biological Imaging Facility and Dr. Tiffany Schmidt for confocal microscope access and the Transgenic and Targeted Mutagenesis Laboratory for cryorecovery of mouse lines. This work was supported by the Beckman Young Investigator Award, William and Bernice E. Bumpus Young Innovator Award, Rita Allen Foundation Scholar Award, and NARSAD Young Investigator and P&S Fund Grant (all Y.K.). M.F.P. was supported by NIH T32 AG20506 and an Arnold O. Beckman Postdoctoral Fellowship.

Received: November 30, 2016

Revised: March 3, 2017

Accepted: May 26, 2017

Published: June 29, 2017

REFERENCES

- Angioni, L., Cocco, C., Ferri, G.L., Argiolas, A., Melis, M.R., and Sanna, F. (2016). Involvement of nigral oxytocin in locomotor activity: a behavioral, immunohistochemical and lesion study in male rats. *Horm. Behav.* **83**, 23–38.
- Bäckman, C.M., Malik, N., Zhang, Y., Shan, L., Grinberg, A., Hoffer, B.J., Westphal, H., and Tomac, A.C. (2006). Characterization of a mouse strain expressing Cre recombinase from the 3' untranslated region of the dopamine transporter locus. *Genesis* **44**, 383–390.
- Banghart, M.R., Neufeld, S.Q., Wong, N.C., and Sabatini, B.L. (2015). Enkephalin disinhibits mu opioid receptor-rich striatal patches via delta opioid receptors. *Neuron* **88**, 1227–1239.
- Bao, S., Chan, V.T., and Merzenich, M.M. (2001). Cortical remodeling induced by activity of ventral tegmental dopamine neurons. *Nature* **412**, 79–83.
- Bayer, S.A., Wills, K.V., Triarhou, L.C., Thomas, J.D., and Ghetti, B. (1995). Systematic differences in time of dopaminergic neuron origin between normal mice and homozygous weaver mutants. *Exp. Brain Res.* **105**, 200–208.
- Beier, K.T., Steinberg, E.E., DeLoach, K.E., Xie, S., Miyamichi, K., Schwarz, L., Gao, X.J., Kremer, E.J., Malenka, R.C., and Luo, L. (2015). Circuit architecture of VTA dopamine neurons revealed by systematic input-output mapping. *Cell* **162**, 622–634.
- Briffaud, V., Williams, P., Courty, J., and Broberger, C. (2015). Excitation of tuberoinfundibular dopamine neurons by oxytocin: crosstalk in the control of lactation. *J. Neurosci.* **35**, 4229–4237.
- Bromberg-Martin, E.S., Matsumoto, M., and Hikosaka, O. (2010). Dopamine in motivational control: rewarding, aversive, and alerting. *Neuron* **68**, 815–834.
- Burkett, J.P., Andari, E., Johnson, Z.V., Curry, D.C., de Waal, F.B.M., and Young, L.J. (2016). Oxytocin-dependent consolation behavior in rodents. *Science* **351**, 375–378.
- Busnelli, M., Saulière, A., Manning, M., Bouvier, M., Galés, C., and Chini, B. (2012). Functional selective oxytocin-derived agonists discriminate between individual G protein family subtypes. *J. Biol. Chem.* **287**, 3617–3629.
- Choe, H.K., Reed, M.D., Benavidez, N., Montgomery, D., Soares, N., Yim, Y.S., and Choi, G.B. (2015). Oxytocin mediates entrainment of sensory stimuli to social cues of opposing valence. *Neuron* **87**, 152–163.
- Crowley, W.R. (2015). Neuroendocrine regulation of lactation and milk production. *Compr. Physiol.* **5**, 255–291.
- Dölen, G., Darvishzadeh, A., Huang, K.W., and Malenka, R.C. (2013). Social reward requires coordinated activity of nucleus accumbens oxytocin and serotonin. *Nature* **501**, 179–184.
- Dubois-Dauphin, M., Barberis, C., and de Bilbao, F. (1996). Vasopressin receptors in the mouse (*Mus musculus*) brain: sex-related expression in the medial preoptic area and hypothalamus. *Brain Res.* **743**, 32–39.
- Duke, D.C., Moran, L.B., Pearce, R.K.B., and Graeber, M.B. (2007). The medial and lateral substantia nigra in Parkinson's disease: mRNA profiles associated with higher brain tissue vulnerability. *Neurogenetics* **8**, 83–94.
- Eliava, M., Melchior, M., Knobloch-Bollmann, H.S., Wahis, J., da Silva Gouveia, M., Tang, Y., Ciobanu, A.C., Triana del Rio, R., Roth, L.C., Althammer, F., et al. (2016). A new population of parvocellular oxytocin neurons controlling magnocellular neuron activity and inflammatory pain processing. *Neuron* **89**, 1291–1304.
- Fujiwara, Y., Hiroyama, M., Sanbe, A., Yamauchi, J., Tsujimoto, G., and Tanoue, A. (2007). Mutual regulation of vasopressin- and oxytocin-induced glucagon secretion in V1b vasopressin receptor knockout mice. *J. Endocrinol.* **192**, 361–369.
- Fuxe, K., Borroto-Escuela, D.O., Romero-Fernandez, W., Ciruela, F., Manger, P., Leo, G., Diaz-Cabiale, Z., and Agnati, L.F. (2012). On the role of volume transmission and receptor-receptor interactions in social behaviour: focus on central catecholamine and oxytocin neurons. *Brain Res.* **1476**, 119–131.
- Gao, S., Becker, B., Luo, L., Geng, Y., Zhao, W., Yin, Y., Hu, J., Gao, Z., Gong, Q., Hurlmann, R., et al. (2016). Oxytocin, the peptide that bonds the sexes also divides them. *Proc. Natl. Acad. Sci. USA* **113**, 7650–7654.
- Gimpl, G., and Fahrenholz, F. (2001). The oxytocin receptor system: structure, function, and regulation. *Physiol. Rev.* **81**, 629–683.
- Grinevich, V., Knobloch-Bollmann, H.S., Eliava, M., Busnelli, M., and Chini, B. (2016). Assembling the puzzle: pathways of oxytocin signaling in the brain. *Biol. Psychiatry* **79**, 155–164.
- Guzman, J.N., Sánchez-Padilla, J., Chan, C.S., and Surmeier, D.J. (2009). Robust pacemaking in substantia nigra dopaminergic neurons. *J. Neurosci.* **29**, 11011–11019.
- Harris, J.A., Hirokawa, K.E., Sorensen, S.A., Gu, H., Mills, M., Ng, L.L., Bohn, P., Mortrud, M., Ouellette, B., Kidney, J., et al. (2014). Anatomical characterization of Cre driver mice for neural circuit mapping and manipulation. *Front. Neural Circuits* **8**, 76.
- Hebb, M.O., and Robertson, H.A. (2000). Identification of a subpopulation of substantia nigra pars compacta gamma-aminobutyric acid neurons that is regulated by basal ganglia activity. *J. Comp. Neurol.* **416**, 30–44.
- Howe, M.W., and Dombeck, D.A. (2016). Rapid signalling in distinct dopaminergic axons during locomotion and reward. *Nature* **535**, 505–510.
- Huber, D., Veinante, P., and Stoop, R. (2005). Vasopressin and oxytocin excite distinct neuronal populations in the central amygdala. *Science* **308**, 245–248.
- Insel, T.R., and Shapiro, L.E. (1992). Oxytocin receptors and maternal behavior. *Ann. N Y Acad. Sci.* **652**, 122–141.

- Insel, T.R., Wang, Z.X., and Ferris, C.F. (1994). Patterns of brain vasopressin receptor distribution associated with social organization in microtine rodents. *J. Neurosci.* *14*, 5381–5392.
- Knobloch, H.S., Charlet, A., Hoffmann, L.C., Eliava, M., Khrulev, S., Cetin, A.H., Osten, P., Schwarz, M.K., Seeburg, P.H., Stoop, R., and Grinevich, V. (2012). Evoked axonal oxytocin release in the central amygdala attenuates fear response. *Neuron* *73*, 553–566.
- Kozorovitskiy, Y., Saunders, A., Johnson, C.A., Lowell, B.B., and Sabatini, B.L. (2012). Recurrent network activity drives striatal synaptogenesis. *Nature* *485*, 646–650.
- Lammel, S., Ion, D.I., Roeper, J., and Malenka, R.C. (2011). Projection-specific modulation of dopamine neuron synapses by aversive and rewarding stimuli. *Neuron* *70*, 855–862.
- Landgraf, R., and Neumann, I.D. (2004). Vasopressin and oxytocin release within the brain: a dynamic concept of multiple and variable modes of neuropeptide communication. *Front. Neuroendocrinol.* *25*, 150–176.
- Lee, B., Yang, C., Chen, T.H., al-Azawi, N., and Hsu, W.H. (1995). Effect of AVP and oxytocin on insulin release: involvement of V1b receptors. *Am. J. Physiol.* *269*, E1095–E1100.
- Lee, H.-J., Caldwell, H.K., Macbeth, A.H., Tolu, S.G., and Young, W.S., 3rd (2008). A conditional knockout mouse line of the oxytocin receptor. *Endocrinology* *149*, 3256–3263.
- Lerner, T.N., Shilyansky, C., Davidson, T.J., Evans, K.E., Beier, K.T., Zalocusky, K.A., Crow, A.K., Malenka, R.C., Luo, L., Tomer, R., and Deisseroth, K. (2015). Intact-brain analyses reveal distinct information carried by SNc dopamine subcircuits. *Cell* *162*, 635–647.
- Liu, Y., and Wang, Z.X. (2003). Nucleus accumbens oxytocin and dopamine interact to regulate pair bond formation in female prairie voles. *Neuroscience* *121*, 537–544.
- Lou, Y., Luo, W., Zhang, G., Tao, C., Chen, P., Zhou, Y., and Xiong, Y. (2014). Ventral tegmental area activation promotes firing precision and strength through circuit inhibition in the primary auditory cortex. *Front. Neural Circuits* *8*, 25.
- Love, T.M. (2014). Oxytocin, motivation and the role of dopamine. *Pharmacol. Biochem. Behav.* *119*, 49–60.
- Ludwig, M., Sabatier, N., Bull, P.M., Landgraf, R., Dayanithi, G., and Leng, G. (2002). Intracellular calcium stores regulate activity-dependent neuropeptide release from dendrites. *Nature* *418*, 85–89.
- Luther, J.A., and Tasker, J.G. (2000). Voltage-gated currents distinguish parvocellular from magnocellular neurones in the rat hypothalamic paraventricular nucleus. *J. Physiol.* *523*, 193–209.
- Manning, M., Stoev, S., Chini, B., Durroux, T., Mouillac, B., and Guillon, G. (2008). Peptide and non-peptide agonists and antagonists for the vasopressin and oxytocin V1a, V1b, V2 and OT receptors: research tools and potential therapeutic agents. *Prog. Brain Res.* *170*, 473–512.
- Marlin, B.J., Mitre, M., D'amour, J.A., Chao, M.V., and Froemke, R.C. (2015). Oxytocin enables maternal behaviour by balancing cortical inhibition. *Nature* *520*, 499–504.
- Mazzuca, M., Minlebaev, M., Shakirzyanova, A., Tyzio, R., Taccola, G., Janackova, S., Gataullina, S., Ben-Ari, Y., Giniatullin, R., and Khazipov, R. (2011). Newborn analgesia mediated by oxytocin during delivery. *Front. Cell. Neurosci.* *5*, 3.
- Mezey, E., and Kiss, J.Z. (1991). Coexpression of vasopressin and oxytocin in hypothalamic supraoptic neurons of lactating rats. *Endocrinology* *129*, 1814–1820.
- Mitre, M., Marlin, B.J., Schiavo, J.K., Morina, E., Norden, S.E., Hackett, T.A., Aoki, C.J., Chao, M.V., and Froemke, R.C. (2016). A distributed network for social cognition enriched for oxytocin receptors. *J. Neurosci.* *36*, 2517–2535.
- Mouillac, B., Chini, B., Balestre, M.N., Elands, J., Trumpp-Kallmeyer, S., Hofflack, J., Hibert, M., Jard, S., and Barberis, C. (1995). The binding site of neuropeptide vasopressin V1a receptor. Evidence for a major localization within transmembrane regions. *J. Biol. Chem.* *270*, 25771–25777.
- Mullis, K., Kay, K., and Williams, D.L. (2013). Oxytocin action in the ventral tegmental area affects sucrose intake. *Brain Res.* *1513*, 85–91.
- Oettl, L.-L., Ravi, N., Schneider, M., Scheller, M.F., Schneider, P., Mitre, M., da Silva Gouveia, M., Froemke, R.C., Chao, M.V., Young, W.S., et al. (2016). Oxytocin enhances social recognition by modulating cortical control of early olfactory processing. *Neuron* *90*, 609–621.
- Ostrowski, N.L., Lolait, S.J., and Young, W.S., 3rd (1994). Cellular localization of vasopressin V1a receptor messenger ribonucleic acid in adult male rat brain, pineal, and brain vasculature. *Endocrinology* *135*, 1511–1528.
- Peris, J., MacFadyen, K., Smith, J.A., de Kloet, A.D., Wang, L., and Krause, E.G. (2017). Oxytocin receptors are expressed on dopamine and glutamate neurons in the mouse ventral tegmental area that project to nucleus accumbens and other mesolimbic targets. *J. Comp. Neurol.* *525*, 1094–1108.
- Pologruto, T.A., Sabatini, B.L., and Svoboda, K. (2003). ScanImage: flexible software for operating laser scanning microscopes. *Biomed. Eng. Online* *2*, 13.
- Purba, J.S., Hofman, M.A., and Swaab, D.F. (1994). Decreased number of oxytocin-immunoreactive neurons in the paraventricular nucleus of the hypothalamus in Parkinson's disease. *Neurology* *44*, 84–89.
- Qiu, F., Qiu, C.-Y., Cai, H., Liu, T.-T., Qu, Z.-W., Yang, Z., Li, J.-D., Zhou, Q.-Y., and Hu, W.-P. (2014). Oxytocin inhibits the activity of acid-sensing ion channels through the vasopressin, V_{1A} receptor in primary sensory neurons. *Br. J. Pharmacol.* *171*, 3065–3076.
- Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis. *Nat. Methods* *9*, 676–682.
- Schorscher-Petcu, A., Sotocinal, S., Ciura, S., Dupré, A., Ritchie, J., Sorge, R.E., Crawley, J.N., Hu, S.-B., Nishimori, K., Young, L.J., et al. (2010). Oxytocin-induced analgesia and scratching are mediated by the vasopressin-1A receptor in the mouse. *J. Neurosci.* *30*, 8274–8284.
- Schultz, W. (2007). Behavioral dopamine signals. *Trends Neurosci.* *30*, 203–210.
- Shah, B.P., Vong, L., Olson, D.P., Koda, S., Krashes, M.J., Ye, C., Yang, Z., Fuller, P.M., Elmquist, J.K., and Lowell, B.B. (2014). MC4R-expressing glutamatergic neurons in the paraventricular hypothalamus regulate feeding and are synaptically connected to the parabrachial nucleus. *Proc. Natl. Acad. Sci. USA* *111*, 13193–13198.
- Shahrokh, D.K., Zhang, T.Y., Diorio, J., Gratton, A., and Meaney, M.J. (2010). Oxytocin-dopamine interactions mediate variations in maternal behavior in the rat. *Endocrinology* *151*, 2276–2286.
- Skuse, D.H., and Gallagher, L. (2009). Dopaminergic-neuropeptide interactions in the social brain. *Trends Cogn. Sci.* *13*, 27–35.
- Song, Z., Borland, J.M., Larkin, T.E., O'Malley, M., and Albers, H.E. (2016). Activation of oxytocin receptors, but not arginine-vasopressin V1a receptors, in the ventral tegmental area of male Syrian hamsters is essential for the reward-like properties of social interactions. *Psychoneuroendocrinology* *74*, 164–172.
- Stoop, R. (2012). Neuromodulation by oxytocin and vasopressin. *Neuron* *76*, 142–159.
- Tang, Y., Chen, Z., Tao, H., Li, C., Zhang, X., Tang, A., and Liu, Y. (2014). Oxytocin activation of neurons in ventral tegmental area and interfascicular nucleus of mouse midbrain. *Neuropharmacology* *77*, 277–284.
- Tribollet, E., Barberis, C., Jard, S., Dubois-Dauphin, M., and Dreifuss, J.J. (1988). Localization and pharmacological characterization of high affinity binding sites for vasopressin and oxytocin in the rat brain by light microscopic autoradiography. *Brain Res.* *442*, 105–118.
- Tyzio, R., Cossart, R., Khalilov, I., Minlebaev, M., Hübner, C.A., Represa, A., Ben-Ari, Y., and Khazipov, R. (2006). Maternal oxytocin triggers a transient inhibitory switch in GABA signaling in the fetal brain during delivery. *Science* *314*, 1788–1792.
- Ungless, M.A., and Grace, A.A. (2012). Are you or aren't you? Challenges associated with physiologically identifying dopamine neurons. *Trends Neurosci.* *35*, 422–430.

- Vaccari, C., Lolait, S.J., and Ostrowski, N.L. (1998). Comparative distribution of vasopressin V1b and oxytocin receptor messenger ribonucleic acids in brain. *Endocrinology* 139, 5015–5033.
- Veinante, P., and Freund-Mercier, M.J. (1997). Distribution of oxytocin- and vasopressin-binding sites in the rat extended amygdala: a histoautoradiographic study. *J. Comp. Neurol.* 383, 305–325.
- Vong, L., Ye, C., Yang, Z., Choi, B., Chua, S., Jr., and Lowell, B.B. (2011). Leptin action on GABAergic neurons prevents obesity and reduces inhibitory tone to POMC neurons. *Neuron* 71, 142–154.
- Walsh, J.J., and Han, M.H. (2014). The heterogeneity of ventral tegmental area neurons: projection functions in a mood-related context. *Neuroscience* 282, 101–108.
- Wanat, M.J., Hopf, F.W., Stuber, G.D., Phillips, P.E.M., and Bonci, A. (2008). Corticotropin-releasing factor increases mouse ventral tegmental area dopamine neuron firing through a protein kinase C-dependent enhancement of I_h . *J. Physiol.* 586, 2157–2170.
- Watabe-Uchida, M., Zhu, L., Ogawa, S.K., Vamanrao, A., and Uchida, N. (2012). Whole-brain mapping of direct inputs to midbrain dopamine neurons. *Neuron* 74, 858–873.
- Wise, R.A. (2004). Dopamine, learning and motivation. *Nat. Rev. Neurosci.* 5, 483–494.
- Young, L.J., and Wang, Z. (2004). The neurobiology of pair bonding. *Nat. Neurosci.* 7, 1048–1054.
- Zheng, J.J., Li, S.J., Zhang, X.D., Miao, W.Y., Zhang, D., Yao, H., and Yu, X. (2014). Oxytocin mediates early experience-dependent cross-modal plasticity in the sensory cortices. *Nat. Neurosci.* 17, 391–399.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit anti-tyrosine hydroxylase	Millipore	Cat#AB152; RRID: AB_390204
Mouse anti-tyrosine hydroxylase	Abcam	Cat#AB129991; RRID: AB_11156128
Rabbit anti-oxytocin	Peninsula Laboratories	Cat#T-4084; RRID: AB_518524
Rabbit anti-oxytocin receptor	Alomone Labs	Cat#AVR-013; RRID: AB_2651123
Goat anti-mouse Alexa 488/594/647	Thermo Fisher Scientific	Cat#A-11029/11032/21236; RRID: AB_2534088; RRID: AB_2534091; RRID: AB_141725
Goat anti-rabbit Alexa 488/594/647	Thermo Fisher Scientific	Cat#A-11034/11037/21244; RRID: AB_2576217, RRID: AB_2534095, RRID: AB_2535812
Rabbit anti-oxytocin receptor (OxtR-2)	Gift from R. Froemke	Marlin et al., 2015 ; Mitre et al., 2016
Chemicals, Peptides, and Recombinant Proteins		
Oxytocin	Tocris	Cat#1910; CAS 50-56-6
L-368,899 hydrochloride	Tocris	Cat#2641; CAS 160312-62-9
L741,626	Tocris	Cat#1003; CAS 81226-60-0
SR 49059	Tocris	Cat#2310; CAS 150375-75-0
CGP54626 hydrochloride	Tocris	Cat#1088; CAS 149184-21-4
NBQX disodium salt	Tocris	Cat#1044; CAS 479347-86-9
(RS)-CPP	Tocris	Cat#0173; CAS 100828-16-8
SR95531 hydrobromide	Tocris	Cat#1262; CAS 104104-50-9
Gallein	Tocris	Cat#3090; CAS 2103-64-2
WAY 267464 dihydrochloride	Tocris	Cat#3933; CAS 1432043-31-6
SSR149415	Axon Medchem	Cat#CS-CM-00328; CAS 439687-69-1
Neurobiotin 488	Vector Laboratories	Cat#SP-1125
Alexa 488 hydrazide	Thermo Fisher Scientific	Cat#A10436
Alexa 594 hydrazide	Thermo Fisher Scientific	Cat#A10438
Green retrobeads (GRBs)	Lumafuor	Cat#G180
Red retrobeads (RRBs)	Lumafuor	Cat#R180
Critical Commercial Assays		
RNAscope Fluorescence Multiplex Assay	ACDBio	N/A
RNAscope Probe- Mm-Slc32a1-C3	ACDBio	Cat#319191-C3
RNAscope Probe- Mm-Th-C2	ACDBio	Cat#317621-C2
RNAscope Probe- Mm-Oxtr	ACDBio	Cat#412171
RNAscope Probe- Mm-Avpr1a	ACDBio	Cat#418061
Experimental Models: Organisms/Strains		
Mouse: C57BL/6	Charles River	Cat#000664; RRID: IMSR_JAX:000664
Mouse: B6.129S-Oxtr ^{tm1.1(cre)Dolsn/J}	Jackson Laboratory	Cat#024234; RRID: IMSR_JAX:024234
Mouse: B6.SJL-Slc6a3 ^{tm1.1(cre)Bkmm/J}	Jackson Laboratory	Cat#006660; RRID: IMSR_JAX:006660
Mouse: Slc32a1 ^{tm2(cre)Lowl} Hze	Jackson Laboratory	Cat#016962; RRID: IMSR_JAX:016962
Mouse: B6.Cg-AVP ^{tm1.1(cre)Hze}	Jackson Laboratory	Cat#023530; RRID: IMSR_JAX:023530
Mouse: B6.Cg-Gt(ROSA)26Sor ^{tm14(CAG-tdTomato)Hze/J}	Jackson Laboratory	Cat#007914; RRID: IMSR_JAX:007914
Mouse: B6.129(SJL)-Oxtr ^{tm1.1Wsy/J}	Jackson Laboratory	Cat#008471; RRID: IMSR_JAX:008471
Sequence-Based Reagents		
AAV9-EF1a-DIO-hChR2(H134R)-mCherry	UPenn Viral Core	CS0543-3CS
AAV9-EF1a-DIO-hChR2(H134R)-eYFP	UPenn Viral Core	CS0633-3CS
AAV9-EF1a-DIO-eYFP	UPenn Viral Core	CS0659-3CS

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Software and Algorithms		
GraphPad Prism 5	GraphPad	RRID: SCR_002798
FIJI	Schindelin et al., 2012	http://fiji.sc/ ; RRID: SCR_002285
MATLAB	MathWorks	RRID: SCR_001622
Igor Pro	WaveMetrics	RRID: SCR_000325

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be fulfilled by the corresponding author Yevgenia Kozorovitskiy (yevgenia.kozorovitskiy@northwestern.edu).

EXPERIMENTAL MODEL AND SUBJECT DETAILS**Mouse Strains and Genotyping**

Animals were handled according to protocols approved by the Northwestern University Animal Care and Use Committee. Weanling and young adult male and female mice (postnatal days 21–50) were used in this study. Approximately equal numbers of males and females were used for every experiment. All mice were group-housed, with standard enrichment procedures, and littermates were randomly assigned to conditions. C57BL/6 mice used for breeding and backcrossing were acquired from Charles River (Wilmington, MA), and all other mouse lines were acquired from the Jackson Laboratory (Bell Harbor, ME). Four internal ribosomal entry site (IRES) Cre recombinase knock-in mouse lines were used in this study. B6.129S-*Oxt*^{tm1.1(cre)Dolsn}/J mice (#024234), which express the enzyme Cre recombinase under control of the oxytocin promoter, are referred to as *Oxt*^{i-Cre} ([Shah et al., 2014](#)); B6.SJL-*Slc6a3*^{tm1.1(cre)Bkmn}/J mice, which express Cre recombinase under control of the dopamine transporter promoter, are referred to as *DAT*^{i-Cre} ([Bäckman et al., 2006](#)); B6.Cg-*Avp*^{tm1.1(cre)Hze} mice (#023530), which express the enzyme Cre recombinase under control of the vasopressin promoter, are referred to as *Avp*^{i-Cre} ([Harris et al., 2014](#)); *Slc32a1*^{tm2(cre)Lowl} mice (#016962), which express Cre recombinase under control of the vesicular GABA transporter promoter, are referred to as *Vgat*^{i-Cre} ([Vong et al., 2011](#)). Heterozygous Cre positive mice of both sexes were used in experiments. For a subset of experiments, Cre positive mice were crossed to a floxed tdTomato reporter strain (Ai14, Jackson Lab, #007914), or a floxed *Oxtr* strain (B6.129(SJL)-*Oxtr*^{tm1.1Wsy}/J, Jackson Lab, #008471) ([Lee et al., 2008](#)). All of the genotyping primers were based on standard protocols available on the Jackson Lab website.

METHOD DETAILS**Stereotaxic Intracranial Injections and Fiber Optic Implants**

Conditional expression of fluorescent proteins or ChR2 in Cre-containing neurons was achieved using recombinant adeno-associated viruses (rAAVs) encoding a double-floxed inverted open reading frame (DIO) of target genes. Mice were transduced with one of the following adeno-associated viruses (AAVs) acquired from the UPenn viral core (Philadelphia, PA): AAV9-EF1a-DIO-hChR2(H134R)-mCherry (1.24×10^{13} GC/ml), AAV9-EF1a-DIO-hChR2(H134R)-eYFP (3.55×10^{13} GC/ml), and AAV9-EF1a-DIO-eYFP (5.55×10^{13} GC/ml).

P25–30 mice were anesthetized with ketamine:xylazine (100:10 mg/kg b.w.), received ketoprofen for analgesia, and were placed on a small animal stereotaxic frame (David Kopf Instruments, Tujunga, CA). rAAVs or green/red retrobeads (50–100 nL for single color retrobead injections, 80 nL each for dual color retrobead injections; Lumafluor, Naples, FL) were delivered unilaterally through a pulled glass pipette at a rate of 100 nL/minute using an UltraMicroPump (World Precision Instruments, Sarasota, FL). Injection coordinates for rAAVs in the PVN were: 1.0 mm posterior to bregma, 0.2 mm lateral, 4.4 mm and 4.7 mm below the pia (0.4–0.5 μ l/location); for retrobeads in the VTA, 2.7 mm posterior to bregma, 0.5 mm lateral, and 4.3 mm below the pia; for retrobeads in the SNc, 2.7 mm posterior to bregma, 1.2 mm lateral, and 4.2 mm below the pia; for retrobeads in the dorsal striatum, 1.5 mm anterior to bregma, 1.6 mm lateral, and 3.0 mm below the pia; and for retrobeads in the ventral striatum, 1.5 mm anterior to bregma, 1.6 mm lateral, and 4.4 mm below the pia. The pipette was held at the injection location for 15 min after retrobead injection and 5–10 min after virus injection. Coordinates were slightly adjusted based on mouse age and size. Mice recovered for 7–9 days following retrograde tracer injections and for over two weeks after rAAV injections.

Tissue Processing, Immunohistochemistry, and Imaging

Mice were deeply anaesthetized with isoflurane and transcardially perfused with 4% paraformaldehyde (PFA) in 0.1 M phosphate buffered saline (PBS). Brains were post-fixed for 1–5 days and washed in PBS, prior to sectioning at 50–60 μ m on a vibratome (Leica

Instruments, Nussloch, Germany). No immunoenhancement was used to increase the signal of virally transduced fluorescent proteins. Sections were pretreated in 0.2% Triton X-100 for an hour at RT, then blocked in 10% bovine serum albumin (BSA, Sigma-Aldrich, ST Louis, MO):PBS with 0.05% Triton X-100 for two hours at RT, and they were then incubated for 24–48 hr at 4°C with primary antibody solution in PBS with 0.2% Triton X-100. On the following day, tissue was rinsed in PBS, reacted with secondary antibody for 2 hr at RT, rinsed again, then mounted onto Superfrost Plus slides (ThermoFisher Scientific, Waltham, MA), dried and coverslipped under ProLong Gold antifade reagent with DAPI (Molecular Probes, Life Technologies, Carlsbad, CA) or under glycerol:TBS (3:1) with Hoechst 33342 (1:1000; ThermoFisher Scientific). For acute slices used in electrophysiology experiments, sections were fixed in 4% PFA overnight, washed in PBS, and processed for immunostaining as described above. Primary antibodies used in the study were rabbit anti-tyrosine hydroxylase (1:1000; AB152, Millipore, Billerica, MA), mouse anti-tyrosine hydroxylase (1:1000; AB129991, Abcam), rabbit anti-oxytocin (1:1000; T-4084, Peninsula Laboratories International, San Carlos, CA), and rabbit anti-oxytocin receptor (1:1000; AVR-013, Alomone Labs, Jerusalem, Israel). Alexa Fluor 488-, 594- and 647-conjugated secondary antibodies against rabbit or mouse (Thermo Fisher Scientific, Waltham, MA) were diluted 1:500. The protocol for the OxtR-2 antibody staining (a generous gift of R. Froemke) was based on published procedures (Mitre et al., 2016). Serum was purified by incubating it against OxtR knockout tissue in 1% bovine serum albumin. Brains were postfixed for 2 hr, transferred to 30% sucrose solution in PBS and stored at 4°C overnight. Then, brains were removed from sucrose solution, embedded in Tissue-Tek O.C.T. compound (VWR), stored overnight at –80°C, and sectioned on a cryostat at 18 μm thickness. Sections were rinsed in PBS, blocked for 2 hr in PBS with 0.2% Triton X-100 and 5% donkey serum, and incubated with OxtR-2 antibody serum at 1:250 dilution. For negative controls, blocking solution without antibody was applied. Following a 2 day-long incubation at 4°C in a humidified chamber, sections were rinsed 3 times in PBS and incubated for 2 hr in Alexa Fluor 647-conjugated goat anti-rabbit antibody (Thermo Fisher Scientific, 1:500). Whole sections were imaged with an Olympus VS120 slide scanning microscope (Olympus Scientific Solutions Americas, Waltham, MA). Confocal images were acquired with a Leica SP5 confocal microscope (Leica Microsystems). Depth-matched z stacks of 1 μm -thick optical sections were analyzed in ImageJ (FIJI) (Schindelin et al., 2012).

Quantitative Fluorescence In Situ Hybridization (FISH)

FISH was conducted following previously published procedures (Banghart et al., 2015). C57BL/6 mice were deeply anesthetized by inhalation of isoflurane and decapitated. Brains were quickly removed and frozen in tissue-freezing medium on a mixture of dry ice and ethanol for 5–15 min prior to storage at –80°C. Brains were subsequently cut on a cryostat (Leica CM1850) into 20 μm -thick sections, adhered to Superfrost Plus slides, and frozen at –80°C. Samples were fixed with 4% PFA in 0.1 M PBS at 4°C for 15 min, processed according to the manufacturer's instructions in the RNAscope Fluorescent Multiplex Assay manual for fresh frozen tissue (Advanced Cell Diagnostics, Newark, CA), and coverslipped with ProLong Gold antifade reagent with DAPI (Molecular Probes). The following probes were used: oxytocin receptor (*Oxtr*), vasopressin type 1a receptor (*Avpr1a*), tyrosine hydroxylase channel 2 (*Th*), and Slc32a1 channel 3 (*Slc32a1/Vgat*). Probes were added to slides in combinations of *Oxtr/Th/Vgat* or *Avpr1a/Th/Vgat*, and the Amp4-b fluorescent amplification reagent was used for all experiments. Sections were subsequently imaged on a Leica SP5 confocal microscope in four channels with a 40x objective lens at a zoom of 1.4 at a resolution of 512 \times 512 pixels with 0.5 μm between adjacent z sections. FISH fluorescent signals were averaged four times to minimize background noise. Images were taken across the entirety of the population of dopaminergic neurons in each brain section.

Image Analysis

Co-localization of oxytocinergic tdTomato⁺ cells with GRB⁺ cells and neurons immunoreactive against oxytocin was determined using confocal images of the PVN in 5–7 sections per brain. Co-localization of PVN oxytocin neurons with other cell types, including GABAergic neurons and vasopressin neurons, was quantified by immunostaining tissue from *Vgat*^{i-Cre}; tdTomato and *Avp*^{i-Cre}; tdTomato mice with anti-oxytocin antibody (2–3 mice for each condition). Images were collected and saved for manual counts by one independent blinded observer. The length of oxytocinergic axons in the VTA and SNc was quantified using confocal images of the VTA and SNc, 4 sections per mouse brain. Axonal length was measured in Z projected imaging using Simple Neurite Tracer in FIJI/ImageJ (Schindelin et al., 2012). To quantify OxtR expression in the VTA and SNc, confocal images of OxtR and TH immunolabeling were acquired under the same imaging parameters and analyzed with ImageJ and MATLAB. Three adjacent z stack slices were combined, and TH signal was filtered with a 5 \times 5 Gaussian filter and thresholded. Thresholds were determined by comparing the manual detection to automated detection for several images from the slice, and the same thresholds were used for the remaining images.

FISH images were analyzed using a custom MATLAB script utilizing imreadBF for file loading and a modified version of Fast 2D peak finder. Briefly, four adjacent z stack slices were combined, for a total of ~2 μm of cells. In general, combining between 2 and 3 μm was optimal to ensure that differences in subcellular localization of RNA transcripts did not result in missed colocalization, while minimizing false positive colocalization driven by signal from a second cell at a different depth in a similar x-y position. All channels were thresholded to remove background signal. Watershed segmentation of the image was performed using the DAPI channel information to localize cell bodies. Puncta of FISH molecules were then counted within established cell boundaries; whether a cell was considered positive for a given marker was determined by setting a transcript-dependent threshold of the number of puncta. This threshold was determined by comparing the count of cells given from a manual count of cells to the histogram of puncta per cell given by the automated program for several images from the slice, and applying that threshold in the remaining images. These

stringent parameters for co-localization and the challenges of quantifying low abundance receptor transcripts likely lead to underestimation of receptor-positive population sizes.

Acute Slice Preparation and Electrophysiology

Coronal brain slice preparation was modified from previously published procedures (Kozorovitskiy et al., 2012). Animals were deeply anesthetized by inhalation of isoflurane, followed by a transcardial perfusion with ice-cold, oxygenated artificial cerebrospinal fluid (ACSF) containing (in mM) 127 NaCl, 2.5 KCl, 25 NaHCO₃, 1.25 NaH₂PO₄, 2.0 CaCl₂, 1.0 MgCl₂, and 25 glucose (osmolarity ~310 mOsm/L). After perfusion, the brain was rapidly removed, and immersed in ice-cold ACSF, equilibrated with 95% O₂/5%CO₂. Tissue was blocked and transferred to a slicing chamber containing ice-cold ACSF, supported by a small block of 4% agar. Bilateral 250 μm-thick slices were cut on a Leica VT1000s in a caudo-rostral direction and transferred into a holding chamber with ACSF, equilibrated with 95%O₂/5%CO₂. Slices were incubated at 34°C for ~30 min prior to electrophysiological recording.

Slices were transferred to a recording chamber perfused with oxygenated ACSF at a rate of 2–4 ml/min and ACSF was warmed to 30–34°C by passing it through a feedback-controlled in-line heater (TC-324C; Warner Instruments, Hamden, CT). Whole-cell and cell-attached recordings were obtained from neurons in the VTA and SNc visualized under infrared DODT contrast video microscopy using patch pipettes of ~2–5 MΩ resistance. Dopaminergic neurons were identified on the basis of tdTomato expression whenever possible, or on the basis of the combination of their electrophysiological and morphological properties (Ungless and Grace, 2012; Wanat et al., 2008). In addition, a large proportion of acute brain slices from *Oxt*^{i-Cre} mice used for electrophysiological recording in optogenetic experiments were immersed in fixative and immunolabeled for tyrosine hydroxylase at the end of the experiment, in order to further validate neuronal identity assessed using electrophysiological and morphological parameters.

For recording spontaneous firing rate and membrane potential, the internal solution consisted of (in mM): 135 K-gluconate, 4 KCl, 10 HEPES, 10 Na-phosphocreatine, 4 MgATP, 0.4 Na₂GTP, and 1 EGTA (pH 7.2, ~295 mOsm/L). For the recordings of spontaneous inhibitory postsynaptic currents (sIPSCs), neurons were held at –70 mV with high chloride internal solution consisting of (in mM): 100 CsCl, 35 CsF, 4 MgCl₂, 10 HEPES, 10 Na-phosphocreatine, 4 MgATP, 0.4 Na₂GTP, and 1 EGTA (pH 7.2, ~295 mOsm/L). Alexa Fluor 488 (10–20 μM) or 594 (10–20 μM) or Neurobiotin (0.1%, Neurobiotin 488 tracer, Vector Laboratories, Burlingame, CA) were added to the internal to visualize cell morphology and confirm cell identity and location. Recordings were made using 700B amplifiers (Axon Instruments, Union City, CA); data were sampled at 10 kHz and filtered at 3 kHz with a MATLAB-based acquisition script (MathWorks, Natick, MA). Series resistance, measured with a 5 mV hyperpolarizing pulse in voltage clamp, was under 20 MΩ and was left uncompensated.

To activate ChR2-expressing cell bodies or fibers of oxytocinergic neurons, 10 ms-long light pulses (470 nm, 50 Hz for 15 s, 5–10 mW/mm²) were delivered at the recording site using whole-field illumination through a 60X water-immersion objective (Olympus, Tokyo, Japan) with a PE300 CoolLED illumination system (CoolLED Ltd., Andover, UK). Epifluorescence illumination was used sparingly to minimize ChR2 activation prior to recording and was never used with a GFP filter cube.

For data analysis of pharmacological flow-in experiments, the mean firing rate of each neuron was calculated every 10 s. Neuronal activity was classified as either enhanced or inhibited by oxytocin when the instantaneous firing rate change in response to oxytocin application was more than 20% above or below the baseline average instantaneous firing rate. Mean firing rates for baseline, oxytocin, and washout periods were calculated from traces recorded 2–3 min before oxytocin application, 3–10 min after oxytocin application, and 2–3 min following wash-out.

For *ex vivo* optogenetic experiments, each recorded neuron was first classified as being activated or inactivated by light stimulation. Instantaneous firing rate and resting membrane potential (bin size, 1 s) were compared before (15 s) and after (20 s) light stimulation. A threshold of 20% over baseline measurement was imposed to classify a neuron as light-responsive, and response duration was the time that neuronal firing rate was modulated by over 20% of baseline. Because a subset of dopaminergic neurons did not show spontaneous activity, even at near physiological temperatures, only resting potential changes were analyzed for these cells.

2-Photon Laser Scanning Imaging

We recorded RB-positive or negative oxytocinergic neurons in the PVN with potassium-based internal solution containing Alexa Fluor 488. After 15–20 min-long recording, cell morphology was visualized using Alexa Fluor 488 (10–20 μM) excited with 910 nm light. The beam of an 80 MHz Ti:Sapphire laser (Mai Tai eHP DS, Newport) was directed by a two-dimensional galvanometer scanning mirror system (HSA Galvo 8315K, Cambridge Technology). Fluorescence emission was collected by two PMTs above and below the sample (H10770P, Hamamatsu) after passing through a dichroic beam splitter (FF670-SDi01-26x38, Semrock) and a bandpass filter (FF02-520/28, Semrock). We used ScanImage 3.8 to control scanning parameters and image acquisition of 1.5 μm-thick z stacks through each recorded neuron (Pologruto et al., 2003). Laser intensity was controlled by a Pockels cell and laser power at the sample plane was 10–15 mW. Area of the soma was measured on z-projections with FIJI/ImageJ (Schindelin et al., 2012).

Pharmacology

Pharmacological agents, except SSR 149415 (Nelivaptan, 10 μM, Axon Medchem, Groningen, Netherlands), were acquired from Tocris (Bristol, UK) or Sigma-Aldrich (St. Louis, MO). Drugs were applied by bath perfusion: *Oxt* (1 μM), WAY 267464 (1 μM), L-368,899 hydrochloride (5 μM), L741,626 (10 μM), SR 49059 (5 μM), CGP54626 hydrochloride (10 μM), gallein (20 μM), SR 95531 hydrobromide (10 μM), NBQX (10 μM), and CPP (10 μM).

QUANTIFICATION AND STATISTICAL ANALYSIS

Offline analyses of electrophysiology were performed using MATLAB (Mathworks, Natick, MA) and Igor Pro (Wavemetrics, Portland, OR). All image analysis was carried out in ImageJ (FIJI, NIH) (Schindelin et al., 2012). Whenever possible, data were analyzed blind to condition. For N sizes, both the number of neurons recorded and the number of animals are provided. Sex and age were balanced across groups. Statistical analyses were done using GraphPad Prism 5 software (GraphPad, LaJolla, CA). Group data are expressed as means \pm SEM. For two group comparisons, statistical significance was determined by two-tailed paired or unpaired Student's t tests or non-parametric analogs, when assumptions for parametric testing were not satisfied. Normality was tested using D'Agostino and Pearson omnibus normality test. For multiple group comparisons, one-way analysis of variance (ANOVA) tests were used for normally distributed data, followed by post hoc analyses. For data that were not normally distributed, non-parametric tests for the appropriate group types were used instead, such as Mann-Whitney. $p < 0.05$ was considered statistically significant.

Neuron, Volume 95

Supplemental Information

**Biased Oxytocinergic Modulation
of Midbrain Dopamine Systems**

Lei Xiao, Michael F. Priest, Jordan Nasenbeny, Ting Lu, and Yevgenia Kozorovitskiy




Figure S1, related to Figure 1. Retrograde and anterograde tracing of oxytocinergic projections

(A) Widefield fluorescence image of a coronal brain section from an *Oxt*^{i-Cre}; Ai14 mouse injected with green retrobeads (GRB) into the VTA (top) or the SNc (bottom). Left: grayscale image of tdTomato fluorescence; middle: grayscale image of transported GRB signal; right: merged color image (green, GRB; red, tdTomato). Retrograde signal is visible in the PVN. LH, lateral hypothalamus.

(B) Epifluorescence images of retrobead injection sites (left) and retrobead signal in the PVN of wildtype mice (right). TH IF, tyrosine hydroxylase immunofluorescence; Oxt IF, oxytocin immunofluorescence.

(C) Epifluorescence images show the distribution of oxytocinergic axons from PVN in the LH (left), central amygdala (CeM) (middle), and ventral striatum (right).

(D) Epifluorescence images depict the distribution of virally transduced PVN oxytocinergic axons in the VTA and SNc.




Figure S2, related to Figure 2. Receptor expression and source identity in the PVN oxytocinergic projection to midbrain dopamine regions

(A) OxtR expression in the VTA of wildtype mice (left) and conditional knockouts of OxtR in dopaminergic neurons (*DAT^{i-Cre}; OxtR^{fl/fl}* mice, cKO) (middle). Right: quantification of the intensity of OxtR distribution in

VTA and SNc of wildtype and cKO mice. n = 6 brain sections from 2 wildtype mice, and 7 sections from 2 cKO mice; two-way repeated measures ANOVA and Bonferroni post hoc test, $p < 0.05$. Error bars reflect SEM.

(B) Confirmation of OxtR expression in VTA (left) and SNc (right) of *DAT*^{i-Cre}; Ai14 mice, using OxtR-2 antibody.

(C) Fluorescence *in situ* hybridization images in the VTA. As a control, the *Oxtr* probe was omitted (right). Tyrosine hydroxylase (*Th*, magenta), vesicular GABA transporter (*Vgat*, orange), and oxytocin receptor (*Oxtr*, cyan).

(D) Same as **(C)** but for the *Avpr1a* probe.

(E) Epifluorescence images demonstrate little co-localization between vasopressinergic tdTomato⁺ neurons (magenta) and oxytocin immunolabeling (green) in the PVN.

(F) Left: confocal images showing the overlap between vasopressinergic tdTomato⁺ neurons (magenta) and oxytocin immunolabeling (green) in the PVN. Right: percentage of neurons positive for vasopressin promoter-driven reporter expression only (magenta), immunolabeling for oxytocin only (green), and double-labeled for both (gray). n = 688 neurons from 2 mice.




Figure S3, related to Figure 4. *Oxytocin differentially alters response properties of VTA and SNc dopamine neurons*

(A) Example epifluorescence images of recorded dopamine neurons filled with Neurobiotin Alexa 488 (green) from *DAT⁺Cre*; Ai14 mice. D-V, dorsal-ventral; M-L, medio-lateral.

(B) Whole-cell recording traces showing membrane potential responses to negative current injections in *DAT⁺* neurons in the VTA (left) and in the SNc (right). Larger negative current injections elicited a characteristic 'sag' in membrane potential of dopamine neurons. ΔV reflects the difference between potential changes in response to hyperpolarizing current injections (-20 versus -100 pA).

(C) Summary graphs of passive and active membrane response properties of *DAT⁺* neurons in the VTA and SNc. Unpaired t test, * $p < 0.05$, $n = 17$ and 18 neurons for VTA and SNc, respectively. Error bars reflect SEM.

(D) Top left: example traces of a single spontaneously generated action potential in a VTA dopamine neuron in baseline condition (gray) and following bath application of $1 \mu\text{M}$ oxytocin (green). Top right: summary graphs for oxytocinergic regulation of action potential parameters in VTA dopamine neurons. Bath application of oxytocin decreased the afterhyperpolarization (AHP) and spike height, but not spike width. Spike width is measured at half-maximal spike amplitude; AHP is the amplitude of action potential undershoot phase; spike height is the peak voltage relative to the undershoot phase amplitude. Paired t test, * $p < 0.05$, $n = 9$ neurons from 7 mice. Bottom: Same as top panel, but in the SNc. Bath application of oxytocin decreased spike width and increased spike amplitude of SNc dopamine neuron action potentials. Paired t test, * $p < 0.05$, $n = 10$ neurons from 9 mice. Error bars reflect SEM.

(E) Left: cell-attached recording traces from VTA *DAT⁺* dopamine neurons in the baseline condition and following application of oxytocin. Right: summary graph of firing rate before and during the application of oxytocin. $\text{FR}_{\text{baseline}}, 1.518 \pm 0.341 \text{ Hz}$; $\text{FR}_{\text{oxytocin}}, 1.725 \pm 0.389 \text{ Hz}$; paired t test, * $p < 0.05$, $n = 10$ neurons from 9 mice. Green, *DAT⁺* neurons that enhanced their firing rate by $>20\%$ in response to oxytocin application. Larger circle marks the example neuron.

(F) Same as (E), but in the SNc. FR_{baseline} , 2.028 ± 0.188 Hz; FR_{oxytocin} , 1.697 ± 0.237 Hz; paired t test, * $p < 0.05$, $n = 6$ neurons from 5 mice. Yellow, DAT⁺ neurons that decreased their firing rate by >20% in response to oxytocin application.




Figure S4, related to Figure 5. *Optical stimulation of oxytocin fibers modulates dopamine neuron activity*

(A) Confocal image of DIO-ChR2-mCherry AAV expression in the PVN of an *Oxt*^{i-Cre} mouse, showing substantial signal overlap between somatic mCherry (magenta) and oxytocin immunofluorescence (green).

(B) Same as **(A)**, but for DIO-ChR2-eYFP AAV, used in a subset of experiments.

(C) Examples of two neurons identified as dopaminergic (left) and non-dopaminergic (right) based on electrophysiological and morphological properties assessed during whole-cell recording with Alexa 594 dye in the internal solution (red). Post hoc immunofluorescent labeling against tyrosine hydroxylase (TH, grayscale) confirms their identity. We carried out post hoc immunolabeling for TH on 28 recorded neurons from 11 mice, correctly assigning DA identity in all cases.

(D) Optical activation of oxytocin fibers in the VTA modestly depolarized membrane potential of dopamine neurons that were not spontaneously active in the acute slice at near physiological temperature. Left: membrane potential of one VTA dopamine neuron, with the light stimulation period marked in blue. Right: summary graph showing membrane potential before, during, and after recovery from light-evoked oxytocin release. One-way repeated measures ANOVA with Tukey post hoc test, * $p < 0.05$, $n = 7$ neurons from 4 mice.

(E) Same as **(D)**, but for the SNc. One-way repeated measures ANOVA with Tukey post hoc tests, ** $p < 0.01$, $n = 16$ neurons from 9 mice.

(F) Current clamp traces from a single SNc dopaminergic neuron with 1pA, 2pA, and 5pA current injections. Bottom middle: summary of firing rate (FR) changes as a function of current injections. One-way repeated measures ANOVA with Tukey post hoc test, * $p < 0.05$, *** $p < 0.0001$, $n = 6$ neurons from 4 mice. Bottom right: correlation between FR and membrane voltage changes. Membrane potential change was calculated based on the injected current and input resistance for each neuron. Red circles mark data points from the neuron used in example traces.

(G) Left: coronal brain section containing the VTA and SNc, showing retro-labeled cells mainly in the VTA when green retrobeads (GRB) were injected into the ventral striatum of an *Oxt*^{i-Cre} mouse (inset). Right:

for GRB injected into dorsal striatum, retro-labeled cells primarily localized to the SNc. Green, GRB; magenta, TH immunostaining; blue, nucleic Hoechst stain.




Figure S5, related to Figure 6. Inhibitory transmission underlies oxytocinergic regulation of SNc and VTA dopamine neurons

(A) Left: current-clamp recording of an SNc dopamine neuron, reversibly silenced by optical activation of oxytocin fibers. Right: another example of an SNc dopamine neuron current-clamp recording, but in the presence of antagonists for AMPA, NMDA, and GABA(A) receptors (NBQX, CPP, and SR95531, respectively).

(B) Summary graphs of optical activation of oxytocin fiber-driven inhibition of spontaneous activity in SNc dopamine neurons. Left: in standard ACSF. Right: in the presence of blockers of fast neurotransmission. Baseline refers to the mean firing rate from a 15 second interval immediately before light stimulation. 'Light' refers to the mean firing rate during 15 seconds of light stimulation. 'After' refers to the mean firing rate during a 20 second interval immediately following light stimulation. Antagonists of AMPA, NMDA and

GABA(A) receptors shortened the duration of oxytocin release-mediated inhibition. One-way repeated measures ANOVA with Tukey post hoc tests, ** $p < 0.01$, $n = 10$ and 18 neurons.

(C) Response duration of optogenetic oxytocin release-evoked modulation of dopamine neurons in the VTA and SNc, in standard ACSF and in the presence of blockers of fast neurotransmission. VTA excitation duration, 14.4 ± 0.7 sec; SNc inhibition duration (ACSF), 59.5 ± 5.5 sec; SNc inhibition duration (blockers), 18.4 ± 0.9 sec. Kruskal-Wallis test, *** $p < 0.001$, $n = 10$ neurons in ACSF, and 18 neurons in the presence of blockers. Error bars reflect SEM.

(D) Summary graph of VTA dopamine neuron firing rate changes in response to optical stimulation, in standard ACSF and following blockade of GABA(A)R and GABA(B)R with SR95531 and CGP54626. Wilcoxon matched-pairs signed rank test, * $p < 0.05$, $n = 6$ from 3 mice.